

MODIS Atmosphere QA Plan *for Collection 061*

Includes:

LMH Flag in "Quality_Assurance_5km" Clarification

Cirrus Flag & High Cloud Flag (06_CT) Clarification

Deep Blue Aerosol Update

Aerosol Over Land Update

Water Vapor and Atmosphere Profile Update

Changes to MOD35 QA Bit Field Documentation

Cloud Optical Properties Update

Introductory Bit String Interpretation Update

CSR Flag Documentation

"Quality_Assurance_1km" Update



Version 11
22 Sep 2021

Paul Hubanks¹

¹ NASA Goddard Space Flight Center, Code 613.2/ADNET, Greenbelt, MD 20071

Table of Contents

1.0	Introduction	1
1.1	QA Approach	2
2.0	Level 2 Runtime QA Flag Structure/Information Content	3
2.1	Types of Runtime QA Flags	3
2.2	Reading and Interpreting Runtime QA Flags	6
Appendix A: Runtime QA Flags of Atmosphere Products		11
	Aerosol (04_L2)	12
	Water Vapor (05_L2)	17
	Cloud (06_L2)	20
	Cloud Optical Properties	20
	Cloud Top Properties	26
	Atmospheric Profile (07_L2)	31
	Cloud Mask (35_L2)	36
	Joint Atmosphere (ATML2)	40
Appendix B: Inventory Metadata of Atmosphere Products		44
	Introduction	45
	ECS (or Core) Inventory Metadata	45
	QA Inventory Metadata	47
	Aerosol (04_L2)	48
	Water Vapor (05_L2)	49
	Cloud (06_L2)	50
	Atmospheric Profile (07_L2)	51
	Cloud Mask (35_L2)	52
	Joint Atmosphere (ATML2)	53
Appendix C: Archive Metadata of Atmosphere Products		55
	Introduction	56
	Aerosol (04_L2)	57
	Water Vapor (05_L2)	58
	Cloud (06_L2)	59
	Atmospheric Profile (07_L2)	60
	Cloud Mask (35_L2)	61
	Joint Atmosphere (ATML2)	62
	Daily Global (08_D3)	63
	Eight Day Global (08_E3)	63
	Monthly Global (08_M3)	63

1.0 Introduction

This document describes the QA (quality assessment or quality assurance) plan for all MODIS Atmosphere products. Topics covered include: quality evaluation approach and scenarios, Level 2 (L2) runtime QA structure and information content, types of L2 runtime QA flags, a primer on reading and interpreting L2 runtime QA flags, detail on how Level 2 (L2) confidence flags impact Level 3 (L3) product statistics, and finally a thorough breakdown of the structure and information content of L2 Runtime QA (in Appendix A).

Also included at the end of this document is information on Inventory and Archive Metadata structures that apply to both L2 and L3 products (in Appendix B and C).

It should be noted that within MODIS Atmosphere, the term “QA” is loosely defined to include a wide variety of flags that detail (1) confidence or quality, (2) retrieval processing path, (3) retrieval status or outcome, (4) retrieval method, (5) data or scene characteristics, and (6) metadata or ancillary input source.

Quality assessment is an important element in the sequential data reduction from Level 0 (L0) raw counts to Level 1B (L1B) calibrated radiance, and continually to Level 2 (L2) orbital swath granules and Level 3 (L3) global gridded products. Level 2 MODIS Atmosphere products, retrieved from MODIS instrument and ancillary data, are grouped into user-friendly Hierarchical Data Format (HDF) files based upon the general categories of aerosol, column water vapor, cloud, atmospheric profiles, cloud mask, and joint atmosphere. Level 3 Atmosphere products, which contain hundreds of 1°x1° global gridded parameters/statistics derived from the L2 products, are grouped into HDF files by temporal period: daily, eight-day, and monthly.

MODIS atmosphere products are grouped as follows (Earth Science Data Type (ESDT) names, used to identify each product by their HDF filename, are provided in parentheses):

Level 2 Products

- Aerosol (04_L2)
- Water Vapor (or Precipitable Water) (05_L2)
- Cloud Properties (06_L2)
- Atmospheric Profiles (07_L2)
- Cloud Mask (35_L2)
- Joint Atmosphere (ATML2)

Level 3 Products

- Daily Global (08_D3)

- Eight-day Global (08_E3)
- Monthly Global (08_M3)

Each Level 2 atmospheric parameter is retrieved at a spatial resolution determined by the sensitivity of retrieval, not necessarily on a single field of view (FOV) basis. Resolutions typically range from 1x1 km to 10x10 km. Level 3 atmospheric parameters are computed at 1x1 degree spatial resolution.

There are two main types of flags described in this document: (1.) Runtime QA Flags and (2.) Metadata (either Inventory or Archive).

Runtime QA Flags (which are only computed and stored for Level 2 products) are designed to convey information on retrieval processing path, input data source, scene characteristics, and the estimated quality of the physical parameters retrieved. In addition, this broad group of flags also includes Cloud Mask QA flags (initially derived at 1x1 km resolution) that may be recomputed at the spatial resolution of the retrieval for the determination of cloudy and clear pixels, land surface type, sunglint, day/night, and snow/ice.

Metadata, on the other hand, are to report high-level granule characteristics and calculated statistics in the broad category of either Inventory or Archive Metadata. Inventory Metadata are searchable, whereas the Archive Metadata are for archival (documentation) only.

1.1 QA Approach

The (automatic) operational approach forms the key to the QA process of MODIS atmosphere products. This encompasses the Runtime QA flags and Metadata produced in each product as well as the operational procedures performed in real time at the GDAAC (Goddard DAAC) or MODAPS (MODIS Data Processing System).

The Runtime QA flags produced in Level 2 products are the central piece of this Quality Assurance (QA) Plan. They are designed not only to report success or failure of criteria being used in retrieval but also to estimate the data quality. These resolution-level QA flags provide much more detailed information and thus provide better information to produce Level 3 global products. The MODIS atmosphere Level 3 products are produced to include various statistical quantities (e.g., mean, standard deviation, histogram and regression). Theoretically, in evaluating the quality of the retrieval, the quality of the inputs should also be considered. Thus, the Level 2 QA flags are to reflect only the product itself, with input sources flagged for later evaluation.

Introductory information on the type and structure of Runtime QA Flags, along with a primer on reading and interpreting QA Flags, can be found in Section 2. The key portion of this document is the detailed tabulation of Runtime QA Flags in Appendix A.

2.0 Level 2 Runtime QA Structure/Information Content

The quality of Level 2 product can be (1) inherited from the L1B radiances, or (2) associated with the retrieval process. The pixel-based L1B validity flags comprising information on dead and saturated detectors, calibration failure, etc., are examined by L2 algorithms for determination of the radiometric status of each pixel. This information can prevent further calculations from being performed if the “valid input data” criteria is not met by the given algorithm. The granule-level L1B QA metadata provides summary information for valid and saturated Earth view observations, and can be useful in screening a granule of data. Details about MODIS L1B QA flags can be found in the MODIS L1B QA plan. The structure and information content of MODIS Atmosphere runtime QA are detailed in the following sections.

It should be noted that Runtime QA flags are only found in Level 2 (L2) Atmosphere products. Level 3 (L3) Atmosphere products contain no Runtime QA flags (bit-string SDS's); however L2 runtime QA flags are used to compute (aggregate and weight) statistics in L3.

2.1 Types of Runtime QA Flags

The SDS run time QA flags are stored based upon product resolution. For convenience, the run time QA flags can be divided into three parts: (1) cloud mask flags, (2) product quality flags, and (3) retrieval processing flags.

Cloud Mask Flags are constructed using the Cloud Mask (35_L2). Typically they are based on the flags from the first byte only, however it may contain additional flags read from subsequent bytes. The first byte of the Cloud Mask is shown below:

Bits	Flag Name	Bit Values	Bit Value Definitions
0	Cloud Mask Status Flag	0	Undetermined
		1	Determined
1-2	Cloud Mask Cloudiness Flag	0	Confident Cloudy (or Fill, if Status Flag = 0)
		1	Probably Cloudy
		2	Probably Clear
		3	Confident Clear
3	Day / Night Flag	0	Night (or Fill, if Status Flag = 0)
		1	Day
4	Sunglint Flag	0	Yes (or Fill, if Status Flag = 0)
		1	No
5	Snow / Ice Flag	0	Yes (or Fill, if Status Flag = 0)
		1	No

6-7	Surface Type Flag	0	Ocean or Deep Lakes and Rivers (or Fill)
		1	Coast or Shallow Lakes and Rivers
		2	Desert
		3	Land

For L2 products at 1×1 km resolution, the first byte of the cloud mask will be written and stored, exactly as detailed above. For coarser resolution products (5×5 or 10×10 km), the information for each QA Flag will be determined by science team members based upon the retrieval method (it may be subsampled or averaged). To avoid duplication in products that are built from more than one retrieval algorithm, cloud mask QA flags will be stored only once in each product (HDF file). The only exception is the Cloud Product (06_L2), where the cloud mask QA flags will be stored at both 1×1 and 5×5 km resolutions.

Product Quality Flags are used to indicate the quality of each parameter retrieved at the product spatial resolution. The first bit indicates usefulness of the parameter, followed by two or three bits for the confidence level. Even though one can store a number from 0 to 7 (corresponding to 8 confidence levels) when three bits are allocated for the Confidence Flag, the Atmosphere L3 production processing software was developed to handle only 4 confidence levels (see table below), which can be adequately stored in only two bits. Usefulness Flags should always be read in tandem with Confidence Flags to differentiate between No Confidence but valid data vs. Fill (missing) data.

An example of the typical format for Product Quality Flags is shown below:

Flag Name	Bit Values	Bit Value Definitions	L3 QA-weighting
<i>Example Parameter</i> Usefulness Flag	0	Not useful	Not used in L3
	1	Useful	Used in L3
<i>Example Parameter</i> Confidence Flag	0	No Confidence (or Fill)	0x
	1	Marginal	1x
	2	Good	2x
	3	Very Good	3x

The MODIS Atmosphere L3 processing software makes use of the L2 Usefulness and Confidence flags by creating L3 QA-weighted mean and standard deviation statistics. The QA weighting is performed by weighting each L2 input pixel by its Confidence flag, so that non-fill *no confidence* data has a weight of 0x, *marginal* data has a weight of 1x, *good* data has a weight of 2x, and *very good* data has a weight of 3x in the statistical computation within the L3 one-degree grid box.

It should be noted that L3 “regular” (non QA-weighted) mean and standard deviation statistics are not weighted by L2 Confidence flags and will **include** all *non-fill* L2 input pixels that have their Quality/Confidence flags set to 0; while the QA-weighted mean and standard deviation statistics will **exclude** these pixels. This technique allows for the creation of L3 (QA-weighted) statistics that can selectively

exclude *no confidence* (or experimental) L2 results.

Finally, users should note that (in addition to regular simple statistics) L2 QA Confidence flags are also ignored in the L3 computation of pixel count, histogram, joint histogram, and regression statistics.

Retrieval Processing Flags are used for miscellaneous purposes. The bit length and information content are determined by the responsible science team member. In general, it may contain:

- Physical, algorithm, climatological constraints
- Atmospheric correction (Rayleigh scattering, gaseous absorption)
- Input resource of ancillary data or MODIS product
- Spectral band and detector status (L1B)
- Occurrence of contamination by thin cirrus

Two common retrieval processing flags are the processing path flag and the input data resource flag. The former indicates the retrieval path, the correction due to Rayleigh scattering and gaseous absorption, or the occurrence of error during retrieval. The latter is to indicate the source of input data sets, such as other MODIS products, ancillary data from other satellites, model assimilated data from GMAO (Goddard Modeling and Assimilation Office), or NCEP (National Centers for Environmental Prediction), or climatology.

2.2 Reading and Interpreting Runtime QA Bit Flags


All Level 2 (L2) MODIS Atmosphere HDF data products contain one or more “bit flag” Scientific Data Set (SDS) arrays. Bit flag SDS names may contain the string “Quality_Assurance” or may have a more descriptive name like “Cloud_Mask”. However, they all have one common attribute – bit flag (or bit string) arrays containing one or more flags stored in particular (fixed) bit positions of the array.

Bit and Byte Array Indexing Convention: “Zero-based”

The convention for indexing arrays varies from language to language. Array indexing in FORTRAN typically starts at 1; and array indexing in C typically starts at 0. However, almost all tools used for bit extraction (in both FORTRAN and C) use an index start convention of 0 for both bits and bytes. In addition, the HDF interface is based in the C (0-based) language. Therefore a 0-based start convention for indexing (numbering) the bits and bytes is always used. In summary, the first byte is always called byte 0, and the first (lowest order) bit is always called bit 0.

HDF Bit Ordering Convention:

When reading bit flags from *bit string* SDSs (Quality Assurance or Cloud Mask arrays, for example), the bits will always be numbered from right (bit index #0) to left (bit index #7). That is, the least significant bit (2^0) is on the right and most significant bit (the sign bit) is on the left. A visual example of this format is offered below.



Numerical Whole-Byte Dumps

Users should note that if one uses the HDF utility, *ncdump*, or other tools to dump full-byte values as decimal numbers (that is, decimal numbers ranging from -128 to 127) from *bit flag* SDS arrays. One simply converts the numbers to 8 bit registers of 0 or 1, where the bit tagged as the first bit (bit index #0) is the lowest order (or least significant) bit (2^0); and the bit tagged as the last bit (bit index #7) is the highest order (or most significant) bit (the sign bit).

To decode individual flags, one queries groups of sequential bits (matching the length of each flag) in the proper bit order from 0 to 7 and interprets the value of those bits (using the same *least significant to most significant* convention) to obtain the correct flag value. An example of this *bit flag* decoding is detailed below.

For example, consider the second byte of the Cloud_Mask_5km SDS:

Scientific Data Set (SDS): "Cloud_Mask_5km" (second byte only)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
C6 Sunlint Flag	2	0	Fill <u>or</u> CTP retrieval fail
		1	No Sunlint & CTP retrieval success
		2	Sunlint & CTP retrieval success
C6 Snow / Ice Flag	2	0	Fill <u>or</u> CTP retrieval fail
		1	No Snow/Ice & CTP retrieval success
		2	Snow/Ice & CTP retrieval success
C6 Surface Type Flag	3	0	Fill <u>or</u> CTP retrieval fail
		1	Ocean, Deep Lakes and Rivers & CTP retr. success
		2	Coast, Shallow Lakes and Rivers & CTP retr. success
		3	Desert & CTP retrieval success
		4	Land & CTP retrieval success
		5	All other valid (non-fill) surface types & CTP retr. succ.
Day / Night Flag	1	0	Night (or Fill, if Status Flag = 0)
		1	Day


Suppose the following flags were set:

- Sunlint Flag = 2 (Sunlint & CTP Retrieval Success)
- Snow/Ice Flag = 1 (No Snow/Ice & CTP Retrieval Success)
- Surface Type Flag = 1 (Ocean & CTP Retrieval Success)
- Day/Night Flag = 1 (Day)

Bit-String Dump

The bits for the example case noted above are shown in the figure below. Flags are then positioned from the right; and least to most significant bits are ordered from right to left for each individual flag.

Sample Bit Flags



Evaluation of signed bit strings


Case 1: Bit strings of length 7 bits or fewer

Evaluating signed bit strings is easy if you are looking at bit strings of length 7 or fewer. In this case, the decimal numbers are always positive (and will range from 0 to a maximum number which depends on the number of bits being read (for example, 3 bits allows for a maximum decimal number of 7)) and will follow the standard interpretation convention shown in the figure below.

Case 2: Bit strings of length 8 bits, but with the sign bit set to 0 (positive numbers)

Evaluating signed bit strings similarly easy if you are looking at bit strings of length 8 and with the most significant bit (the sign bit) set to 0. In this case the numbers are (still) always positive (and will range from 0 to 127) and will follow the standard interpretation convention shown in the figure below.

Single Byte Representation



Case 3: Bit strings of length 8 bits, but with the sign bit set to 1 (negative numbers)

Evaluating signed bit strings is more convoluted if you are looking at bit strings of length 8 but with the most significant bit (the sign bit) set to 1. In this case the numbers are always negative (and will range from -1 to -128) and follow the interpretation formula outlined below:

Step 1: With the sign bit set to 1, the number will be given a negative sign.

Step 2: Flip all 8 bit registers. All 0's change to 1's; and all 1's change to 0's.

Step 3: Evaluate the bit registers using the Case 2 convention shown above (shown in the previous figure), however when you are done with that calculation, add 1 to the result. Note that the sign bit will always get flipped back to 0 in step 2.

So in the example shown above, with the bit registers showing 10010110. Step 1 says you will assign a negative (-) sign to the result. Step 2 says you flip all the bit registers, which yields 01101001. This "flipped" bit string is interpreted as:

$$1 \times 2^6 + 1 \times 2^5 + 1 \times 2^3 + 1 \times 2^0$$

Or

$$64 + 32 + 8 + 1$$

Or

$$105$$

Adding 1 (as specified in Step 3) we get,

$$106$$

Changing the sign to negative (as noted in Step 1) we get the final answer

$$-106$$

So the bit string of 10010110, will be read as -106.

Numerical Whole-Byte Dump

Sometimes, it's helpful to dump out the numeric values of byte arrays in order to see what values are being stored/read (as a sanity check) to see if you understand how these flags are being set and if they make sense given the sample granule you are studying. There are several ways to dump out the numeric value of a byte from an HDF bit

flag array. One of the easiest ways is to use the HDF dump utility, *ncdump*. Here is example syntax of using *ncdump* to dump out the byte values of the SDS *Cloud_Mask_5km* from a sample HDF file called *test.hdf* to an output file called *dump.out* on a linux/irix command line:

```
ncdump -v Cloud_Mask_5km test.hdf > dump.out
```

Online Bit-String Conversion Tools

There are online tools (calculators) that can assist users to convert bit strings to decimal values (and visa-versa). Here is one example online tool, the link is below along with a screen capture:

http://www.binaryconvert.com/convert_signed_char.html

The screenshot shows a web-based conversion tool for signed characters. At the top, a navigation bar contains tabs for 'Unsigned char', 'Signed char', 'Unsigned short', 'Signed short', 'Unsigned int', 'Signed int', 'Float', and 'Double'. The 'Signed char' tab is selected, and a blue banner below it reads 'Signed char (8-bit) Two's complement'. The interface is divided into two main sections: 'Decimal' and 'Binary'. The 'Decimal' section, highlighted with a blue button, displays the value '-106' and notes 'accurate representation = -106'. To the right of this section is a large blue sphere icon and a 'New conversion' button. The 'Binary' section, also highlighted with a blue button, displays the bit string '1 0 0 1 0 1 1 0' in a light gray box. Below the bit string are eight small square indicators: the first, third, fifth, and seventh squares are green, while the second, fourth, sixth, and eighth squares are gray.

Appendix A:

Runtime QA Flags

of

Atmosphere Products

Aerosol Product: MOD04_L2 (Terra) & MYD04_L2 (Aqua)

For the Aerosol product, the Runtime QA flags are stored in two Scientific Data Sets (SDSs): *Quality_Assurance_Land* and *Quality_Assurance_Ocean*

Quality_Assurance_Land and *Quality_Assurance_Ocean* are five byte SDSs that contain product quality flags, retrieval processing flags, and input data resource flags designed separately for land and ocean because of differences in the retrieval algorithms. All Aerosol QA Flag arrays have the following characteristics:

Scientific Data Set (SDS) Name: <i>"Quality_Assurance_Land"</i> Description: <i>Product quality and retrieval processing flags over Land</i> Length: <i>6 bytes (48 bits)</i>			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
0.47 μm Aerosol Optical Thickness Usefulness Flag	1	0 1	Not useful (All Products are Fill Values) Useful (Valid Products)
0.47 μm Aerosol Optical Thickness Confidence Flag	3	0 1 2 3	No Confidence (or Fill) Marginal Good Very Good
0.66 μm Aerosol Optical Thickness Usefulness Flag	1	0 1	Not useful Useful
0.66 μm Aerosol Optical Thickness Confidence Flag	3	0 1 2 3	No Confidence (or Fill) Marginal Good Very Good
<i>processing path flags</i>			
Part I: Retrieving condition flags when inversion is performed. Retrieved values are output.	4	0 1 2 3 4 5 6 7 8 9 10 11	Retrieval performed normally (no issues) Procedure 2 performed (semi-bright surface) Water pixels in 10 x10 box Cirrus present Fitting error > 0.25 -0.1 < Retrieved τ < 0.0 Number of pixels between 12 & 20 Number of pixels between 21 & 30 Number of pixels between 31 & 50 Ångstrom out of bounds Retrieved T < 0.2 No Retrieval

Part II: Retrieving condition flags when inversion is NOT performed. Fill values are output.	4	0 1 2 3 4 5 6	No error Solar/sensor geometry out of bounds in LUT Apparent reflectance out of bounds in LUT Number of pixels < 12 $\rho_{2.1 \mu m} > 0.35$ (too bright) Retrieved T < -0.1 Retrieved T > 5.0
Spares	4		TBD
<i>input data resource flags</i>			
Total ozone	2	0 1 2 3	TOVS TOMS Climatology DAO
Total Precipitable Water	2	0 1 2 3	NCEP / GDAS MOD05* NIR (MODIS Near-IR Water Vapor Retrieval) Climatology DAO
Snow Cover	2	0 1	MOD35* (MODIS L2 Cloud Mask) MOD10* (MODIS L3 Eight-day Snow Cover)
Spares	6		TBD
Deep Blue Aerosol Usefulness Flag §	1	0 1	Not useful Useful
Deep Blue Aerosol Confidence Flag §	2	0 1 2 3	No Confidence (or Fill) Marginal Good Very Good
Deep Blue Aerosol Type § <i>Note: Flags 2 and 3 are reversed from the Aerosol Type (over land only)</i>	2	0 1 2 3	Mixed Dust Smoke Sulfate
Spares	3		TBD
Combined Deep Blue & Dark Target 550 Aerosol Usefulness Flag §	1	0 1	Not useful Useful
Combined Deep Blue & Dark Target 550 Aerosol Confidence Flag §	2	0 1 2 3	No Confidence (or Fill) Marginal Good Very Good
Spares	5		TBD

* MOD prefix denotes MODIS/Terra (AM overpass) data -- MYD prefix should be substituted for MODIS/Aqua (PM overpass) data

§ Implemented in PGE04 versions 5.3.0 and later (Initially implemented in the Collection 005 Aqua reprocess)

Scientific Data Set (SDS): “Quality_Assurance_Ocean” Description: Product quality and retrieval processing flags over Ocean Length: 5 bytes (40 bits)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Aerosol Parameters (Best Soln.) Usefulness Flag	1	0 1	Not useful Useful
Aerosol Parameters (Best Soln.) Confidence Flag	3	0 1 2 3	No Confidence (or Fill) Marginal Good Very Good
Aerosol Parameters (Avg. Soln.) Usefulness Flag	1	0 1	Not useful Useful
Aerosol Parameters (Avg. Soln.) Confidence Flag	3	0 1 2 3	No Confidence (or Fill) Marginal Good Very Good
<i>processing path flags</i>			
Part I: Aerosol over Ocean Retrieving Condition when inversion is NOT performed. <i>Note: τ (550 nm) fill value will be output on conditions 1 through 10.</i>	4	0 1 2 3 4 5 6 7 8 9 10	Retrieval is performed Glitter is present Cloudy R (0.865 μm) too low for retrieving optical thickness Total number of available VIS / SWIR wavelength bands (from 550 to 1240 nm) is insufficient Total number of available wavelengths < 3 Angles Out-of-Bounds Land present in 10 x 10 km box τ (550 nm) < -0.01; algorithm found negative values of optical thickness (there is a problem) τ (550 nm) > 5.0; out of bounds in lookup table All Channels do not have valid data

Part II: Aerosol over Ocean Retrieving Condition when inversion IS performed. <i>Note: τ (550 nm) retrieved value will be output on conditions 0 through 10.</i> <i>Confidence Flag Notes:</i> <ul style="list-style-type: none"> • If Retrieving Condition = 0, then Confidence Flag is set to Very Good (3). • If = 7 or 14 then Confidence Flag is set to Good (2). • If = 1, 3, 4, 6, 8 or 10, then Confidence Flag is set to Marginal (1). • If = 2, 5, 9, 11, 12, 13, 15 then Confidence Flag is set to No Confidence (0). 	4	0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15	Retrieval performed normally Number of useful pixels within 10 x 10 km box is < 10% R (0.865 μm) low but large enough for retrieving optical thickness; the size distribution is questionable; η = fill value 1.65 μm channel not used 2.13 μm channel not used 1.65 and 2.13 μm channels not used Large uncertainty in both retrieved τ and aerosol type Large uncertainty in retrieved τ , but aerosol type is stable. The best value of ϵ is larger than the threshold value (5%) $-0.01 < \tau$ (550 nm) < 0 but to avoid bias in level 3 product Glint angle between 30° and 40° Glint: store only reflectance, SD and Number of pixels used Glint thick dust retrievals Possible cirrus contamination Off glint thick dust retrievals No retrieval performed
<i>input data resource flags</i>			
Total Ozone	2	0 1 2 3	TOMS TOVS Climatology DAO
Total Precipitable Water	2	0 1 2 3	NCEP/GDAS MOD05* NIR (MODIS Near-IR Water Vapor Retrieval) Climatology DAO
Snow Cover	2	0 1	MOD35* (MODIS Cloud Mask) MOD10* (MODIS Eight-day Snow Cover)
Spares	2		TBD
Spares	8		TBD
Spares	8		TBD

* MOD prefix denotes MODIS/Terra (AM overpass) data -- MYD prefix should be substituted for MODIS/Aqua (PM overpass) data

Sample Runtime QA Flag Images from the Aerosol Product


Reference L1B Image of Scene

12 Feb 2001 15:10



MODIS Terra (AM)


R:G:B = Band 1 : Band 4 : Band 3

Flag = "Aerosol Optical Thickness (0.47 μm) Usefulness"

SDS = "Quality_Assurance_Land"

(Aerosol over Land)


Flag = "Surface Type Flag"



SDS = "Cloud_Mask_QA"

(Cloud Mask)


Flag = "Aerosol Type Flag (over Land only)"



SDS = "Quality_Assurance_Land"

(Aerosol over Land)


Flag = "Sunlint Flag"



SDS = "Cloud_Mask_QA"

(Cloud Mask)

Flag = "Part I: Aerosol over Ocean Retrieving Condition"



SDS = "Quality_Assurance_Ocean"

(Aerosol over Ocean)

Note: Images were created by "bitflag_visualizer", available at http://modis-atmos.gsfc.nasa.gov/tools_bitflag_visualizer.html.

Water Vapor Product: MOD05_L2 (Terra) & MYD05_L2 (Aqua)

The Water Vapor (or Total Precipitable Water) product combines results from both the NIR (1×1 km) and IR algorithms (5 × 5 km). For near infrared (NIR) total precipitable water, the first byte contains cloud mask QA (1 × 1 km), and the second byte (a separate array from the cloud mask QA) contains NIR product quality and retrieval processing flags. For the five bytes of IR total precipitable water results, only product quality and retrieval processing flags are stored. Since IR total precipitable water results are copied from 07_L2, the cloud mask related QA flags can be retrieved from 07_L2, and therefore it will not be duplicated here in the 05_L2 product. All Water Vapor QA Flag arrays have the following characteristics:

- Spatial resolution: 1 × 1 km (NIR) and 5 × 5 km (IR)
- Processing mode: Daytime only (NIR) and Both Day and Night (IR)

<i>Scientific Data Set (SDS): "Cloud_Mask_QA"</i> <i>Description: Cloud mask QA flags at 1x1 km</i> <i>Length: 1 byte (8 bits)</i>			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Mask Status Flag	1	0	Undetermined
		1	Determined
Cloud Mask Cloudiness Flag	2	0	Confident Cloudy (or Fill, if Status Flag = 0)
		1	Probably Cloudy
		2	Probably Clear
		3	Confident Clear
Day / Night Flag	1	0	Night (or Fill, if Status Flag = 0)
		1	Day
Sunglint Flag	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Snow / Ice Flag	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Surface Type Flag	2	0	Ocean or Deep Lakes and Rivers (or Fill)
		1	Coast or Shallow Lakes and Rivers
		2	Desert
		3	Land

Scientific Data Set (SDS): “Quality_Assurance_Near_Infrared” Description: Product quality and retrieval processing flags for Precipitable Water (Near IR) at 1x1 km Length: 1 byte (8 bits)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Total Precipitable Water (NIR) Usefulness Flag	1	0 1	Not useful Useful
Total Precipitable Water (NIR) Confidence Flag	3	0 1 2 3	Bad (or Fill) Marginal Confidence Good Confidence Very Good Confidence
Inversion Method Used (NIR)	2	0 1 2	Two channel ratio Three channel ratio No retrieval
Surface Type	2	0 1 2 3	Bright Land Clear Ocean Cloud Sunglint

Scientific Data Set (SDS): “Quality_Assurance_Infrared” Description: Product quality and retrieval processing flags for Precipitable Water (IR) at 5x5 km Length: 5 bytes (40 bits)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Total Precipitable Water (IR) Usefulness Flag ⁿ	1	0 1	Not useful Useful
Total Precipitable Water (IR) Confidence Flag ⁿ	3	0 1 2 3	Bad (or Fill) Best Quality <i>not currently used</i> <i>not currently used</i>
Spares	4		TBD
Number of Cloudy Pixels	Int 8	0-25	Number of 1 km Cloudy pixels in 5x5 retrieval area
Number of Clear Pixels	Int 8	0-25	Number of 1 km Clear pixels in 5x5 retrieval area
Number of Missing Pixels	Int 8	0-25	Number of 1 km Missing pixels in 5x5 retrieval area
IR Water Vapor Retrieval Method Used	2	0 1 2 3	Split Window (11-12 μ m) Technique Integration of Moisture Profile Other No Retrieval
Spares	6		TBD

ⁿ Flag is not reliable in Not Useful / Bad regions (i.e. Not Useful pixels are incorrectly tagged as Useful, etc. about ½ the time.)

Sample Runtime QA Flag Images from the Water Vapor Product

Reference L1B Image of Scene


12 Feb 2001 15:10



MODIS Terra (AM)


R:G:B = Band 1 : Band 4 : Band 3

Flag = "Surface Type Flag (NIR Retrieval)"




SDS = "Quality_Assurance_Near_Infrared" (NIR Water Vapor)

Flag = "Cloud Mask Cloudiness Flag"



SDS = "Cloud_Mask_QA"

(Cloud Mask)

Flag = "Total Precipitable Water (IR) Usefulness Flag" ¹

SDS = "Quality_Assurance_Infrared"

(IR Water Vapor)

¹ TPWIR Flags are copied from the 07_L2 Atmospheric Profile product -- and all 07_L2 Atmospheric Profile Usefulness and Confidence Flags are noisy.

Note: Images were created by "bitflag_visualizer", available at http://modis-atmos.gsfc.nasa.gov/tools_bitflag_visualizer.html.

Cloud Product: MOD06_L2 (Terra) & MYD06_L2 (Aqua)

The MODIS Cloud product consists of a 1 km set of parameters derived from solar reflectance channels (Cloud Optical Properties and Cirrus Reflectance) and a 5 km set of parameters determined from thermal emitted channels (Cloud Top Properties).

Cloud Optical Properties

Cloud Optical Property QA flags are stored in 2 separate QA arrays (SDS's). The first SDS, *Cloud_Mask_1km*, contains Cloud Mask QA flags, which are copied from the 35_L2 Cloud Mask product. The second SDS (*Quality_Assurance_1km*) contains product quality, retrieval processing, and scene characteristic flags. Detail is provided below.

- Spatial resolution: 1 x 1 km
- Processing mode: Daytime only

<i>Scientific Data Set (SDS): "Cloud_Mask_1km"</i> <i>Description: Cloud mask QA flags at 1x1 km</i> <i>Length: 2 bytes (16 bits)</i>			
Flag Name	Number of Bits	Bit Value	Bit Value Definitions
Cloud Mask Status Flag	1	0	Undetermined ^{n,f}
		1	Determined
Cloud Mask Cloudiness Flag	2	0	Confident Cloudy (or Fill, if Status Flag = 0)
		1	Probably Cloudy
		2	Probably Clear
		3	Confident Clear
Day / Night Flag	1	0	Night ^{n,f} (or Fill, if Status Flag = 0)
		1	Day
Sunglint Flag	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Snow / Ice Flag	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Surface Type Flag	2	0	Ocean or Deep Lakes and Rivers (or Fill)
		1	Coast or Shallow Lakes and Rivers
		2	Desert
		3	Land
Heavy Aerosol Flag	1	0	Yes ^{n,f} (or Fill, if Status Flag = 0)
		1	No
Thin Cirrus Flag <i>(Based on low threshold using 1.38 μm band.)</i>	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Shadow Flag	1	0	Yes ^{n,f} (or Fill, if Status Flag = 0)
		1	No
Spares	5		TBD

Note that the *Quality_Assurance_1km* SDS in 06_L2 HDF files was expanded from 5 bytes (in Collection 005/051) to 9 bytes (in Collection 006).

Scientific Data Set (SDS): <i>“Quality_Assurance_1km”</i>			
Description: <i>Cloud Optical Property product quality and retrieval processing QA flags at 1x1 km</i>			
Length: <i>C006 = 9 bytes (72 bits)</i> <i>C005/051 = 5 bytes (40 bits)</i>			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Primary (VNSWIR - 2.1 μm) Cloud Optical Thickness Usefulness Flag	1	0 1	Not useful Useful
Primary (VNSWIR - 2.1 μm) Cloud Optical Thickness Confidence Flag <i>(This flag is not important in C6)</i>	2	0 1 2 3	No Confidence or Fill <i>(Fill setting is used if both Cloudy COT and Partly Cloudy (PCL) COT are fill)</i> Marginal Confidence <i>(Marginal setting not used in C6)</i> Good Confidence <i>(Good setting not used in C6)</i> Very Good Confidence <i>(Very Good setting is default for all non-fill retrievals, however the QA might be associated with either a non-fill Cloudy COT or a non-fill Partly Cloudy (PCL) COT, the other might be fill)</i>
Surface Type Flag <i>(New Flag for Collection 061)</i> <i>(Previously Spares in C6 and earlier)</i>	2	0 1 2 3	Ice-Free Ocean (or Fill) Ice Covered Ocean Snow-Free Land Snow Covered Land
Primary (VNSWIR - 2.1 μm) Cloud Effective Radius Usefulness Flag	1	0 1	Not useful Useful
Primary (VNSWIR - 2.1 μm) Cloud Effective Radius Confidence Flag <i>(This flag is not important in C6)</i>	2	0 1 2 3	No Confidence or Fill <i>(Fill setting is used if both Cloudy CER and Partly Cloudy (PCL) CER are fill)</i> Marginal Confidence <i>(Marginal setting not used in C6)</i> Good Confidence <i>(Good setting not used in C6)</i> Very Good Confidence <i>(Very Good setting is default for all non-fill retrievals, however the QA might be associated with either a non-fill Cloudy CER or a non-fill Partly Cloudy (PCL) CER, the other might be fill)</i>
Primary (VNSWIR - 2.1 μm) Cloud Water Path Usefulness Flag	1	0 1	Not useful Useful
Primary (VNSWIR - 2.1 μm) Cloud Water Path Confidence Flag <i>(This flag is not important in C6)</i>	2	0 1 2 3	No Confidence or Fill <i>(Fill setting is used if both Cloudy CWP and Partly Cloudy (PCL) CWP are fill)</i> Marginal Confidence <i>(Marginal setting not used in C6)</i> Good Confidence <i>(Good setting not used in C6)</i> Very Good Confidence <i>(Very Good setting is default for all non-fill retrievals, however the QA might be associated with either a non-fill Cloudy CWP or a non-fill Partly Cloudy (PCL) CWP, the other might be fill)</i>

Cloud Retrieval Phase Flag <i>(Cloud Retrieval Phase Flag duplicated from the 3rd byte). For combining with the 1.6 - 2.1 μm Cloud Retrieval Outcome Flag (below). Needed by L3 to properly compute 1621 Cloud Fractions. (See page 25 of this Plan)</i>	3	0 1 2 3 4	Cloud Mask Undetermined or Non-Snow Land ^{n,f} Not Processed (typically clear) ^f Liquid Water Cloud Ice Cloud Undetermined Phase Cloud
1.6 - 2.1 μ m Cloud Retrieval Outcome Flag <i>The Cloud Retrieval Phase Flag and 1621 Outcome Flag are read as a combined flag by L3 to properly compute 1621 Cloud Retrieval Fractions. (See page 25 of this Plan)</i>	1	0 1	Retrieval not attempted or unsuccessful ^f Retrieval successful <i>(over ocean, snow, & ice only)</i>
Spare	1		TBD
Cloud Retrieval Phase Flag <i>Primary Cloud Retrieval Phase Flag and Outcome Flag are read as a combined flag by L3 to properly compute Primary Cloud Retrieval Fractions. (See page 25 of this Plan)</i>	3	0 1 2 3 4	Cloud Mask Undetermined ^{n,f} Not Processed (typically clear) ^f Liquid Water Cloud Ice Cloud Undetermined Phase Cloud
Primary (VNSWIR - 2.1 μ m) Cloud Retrieval Outcome Flag <i>Primary Cloud Retrieval Phase Flag and Outcome Flag are read as a combined flag by L3 to properly compute Primary Cloud Retrieval Fractions. (See page 25 of this Plan)</i>	1	0 1	Retrieval not attempted or unsuccessful ^f Retrieval successful
Rayleigh Correction	1	0 1	No Yes, correction was made
Atmospheric Water Vapor Correction	1	0 1	No Yes, correction was made
Band Used for Primary Optical Thickness Retrieval	2	0 1 2 3	Retrieval not attempted ^f 0.645 μ m (land) 0.858 μ m (water) 1.24 μ m (snow / ice)
1.6 - 2.1 μ m Cloud Optical Thickness Usefulness Flag	1	0 1	Not useful Useful
1.6 - 2.1 μ m Cloud Optical Thickness Confidence Flag <i>(This flag is not important in C6)</i>	2	0 1 2 3	No Confidence or Fill <i>(Fill setting is used if both Cloudy COT and Partly Cloudy (PCL) COT are fill)</i> Marginal Confidence <i>(Marginal setting not used in C6)</i> Good Confidence <i>(Good setting not used in C6)</i> Very Good Confidence <i>(Very Good setting is default for all non-fill retrievals, however the QA might be associated with either a non-fill Cloudy COT or a non-fill Partly Cloudy (PCL) COT, the other might be fill)</i>
1.6 - 2.1 μ m Cloud Effective Radius Usefulness Flag	1	0 1	Not useful Useful

1.6 - 2.1 μm Cloud Effective Radius Confidence Flag <i>(This flag is not important in C6)</i>	2	0 1 2 3	No Confidence or Fill <i>(Fill setting is used if both Cloudy CER and Partly Cloudy (PCL) CER are fill)</i> Marginal Confidence <i>(Marginal setting not used in C6)</i> Good Confidence <i>(Good setting not used in C6)</i> Very Good Confidence <i>(Very Good setting is default for all non-fill retrievals, however the QA might be associated with either a non-fill Cloudy CER or a non-fill Partly Cloudy (PCL) CER, the other might be fill)</i>
Clear Sky Restoral Type Flag	2	0 1 2 3	Not Restored Tagged as "Partly Cloudy" (PCL) via Edge Detection Restored to Clear Sky via Spatial Variance Tagged as "Partly Cloudy" (PCL) via 250 meter Tests
1.6 - 2.1 μm Cloud Water Path Usefulness Flag	1	0 1	Not useful Useful
1.6 - 2.1 μm Cloud Water Path Confidence Flag <i>(This flag is not important in C6)</i>	2	0 1 2 3	No Confidence or Fill <i>(Fill setting is used if both Cloudy CWP and Partly Cloudy (PCL) CWP are fill)</i> Marginal Confidence <i>(Marginal setting not used in C6)</i> Good Confidence <i>(Good setting not used in C6)</i> Very Good Confidence <i>(Very Good setting is default for all non-fill retrievals, however the QA might be associated with either a non-fill Cloudy CWP or a non-fill Partly Cloudy (PCL) CWP, the other might be fill)</i>
Primary Cloud Retrieval (VNSWIR - 2.1 μm) Multilayer Cloud & Phase Flag	3	0 1 2 3 4 5 6 7	Cloud mask undetermined ^{n,f} Not Processed (typically clear) ^f Single-Layer Liquid Water Cloud Multi-Layer Liquid Water Cloud Single-Layer Ice Cloud Multi-Layer Ice Cloud Single-Layer Undetermined Phase Cloud Multi-Layer Undetermined Phase Cloud
Primary Cloud Retrieval (VNSWIR - 2.1 μm) Outcome Flag <i>(Primary Cloud Retrieval Outcome Flag duplicated from the 3rd byte). For combining with the Primary Cloud Retrieval Multilayer Cloud & Phase Flag (above). Needed by L3 to properly compute 1L & ML Cloud Fractions.</i>	1	0 1	Retrieval not attempted or unsuccessful ^f Retrieval successful
Spare	1		TBD
Phase Difference Multilayer Test	1	0 1	No Yes
Delta Precipitable Water Multilayer Test	1	0 1	No Yes
Delta Precipitable Water at 900mb Test	1	0 1	No Yes

Tau Difference VIS-NIR Multilayer Test	1	0 1	No Yes
Pavlonis-Heidinger Multilayer Test	1	0 1	No Yes
Spares	3		TBD
VNSWIR - 1.6 μm Cloud Retrieval Phase & Outcome <i>(The Cloud Retrieval Phase Flag and Outcome Flag can be read as a "combined" flag as documented here -- or read as separate flags -- the bit structure is identical. (See page 25 of this Plan for details))</i>	4	0 1 2 3 4 10 11 12	Cloud Mask Undetermined ^{n,f} Not Processed (typically clear) ^f Failed Liquid Water Cloud Retrieval Failed Ice Cloud Retrieval Failed Undetermined Phase Cloud Retrieval Successful Liquid Water Cloud Retrieval Successful Ice Cloud Retrieval Successful Undetermined Phase Cloud Retrieval
VNSWIR - 1.6 μm PCL (Partly Cloudy) Cloud Retrieval Phase & Outcome <i>(The Cloud Retrieval Phase Flag and Outcome Flag can be read as a "combined" flag as documented here -- or read as separate flags -- the bit structure is identical. (See page 25 of this Plan for details))</i>	4	0 1 2 3 4 10 11 12	Cloud Mask Undetermined ^{n,f} Not Processed (typically clear) ^f Failed Liquid Water Cloud Retrieval Failed Ice Cloud Retrieval Failed Undetermined Phase Cloud Retrieval Successful Liquid Water Cloud Retrieval Successful Ice Cloud Retrieval Successful Undetermined Phase Cloud Retrieval
VNSWIR - 3.7 μm Cloud Retrieval Phase & Outcome <i>(The Cloud Retrieval Phase Flag and Outcome Flag can be read as a "combined" flag as documented here -- or read as separate flags -- the bit structure is identical. (See page 25 of this Plan for details))</i>	4	0 1 2 3 4 10 11 12	Cloud Mask Undetermined ^{n,f} Not Processed (typically clear) ^f Failed Liquid Water Cloud Retrieval Failed Ice Cloud Retrieval Failed Undetermined Phase Cloud Retrieval Successful Liquid Water Cloud Retrieval Successful Ice Cloud Retrieval Successful Undetermined Phase Cloud Retrieval
VNSWIR - 3.7 μm PCL (Partly Cloudy) Cloud Retrieval Phase & Outcome <i>(The Cloud Retrieval Phase Flag and Outcome Flag can be read as a "combined" flag as documented here -- or read as separate flags -- the bit structure is identical. (See page 25 of this Plan for details))</i>	4	0 1 2 3 4 10 11 12	Cloud Mask Undetermined ^{n,f} Not Processed (typically clear) ^f Failed Liquid Water Cloud Retrieval Failed Ice Cloud Retrieval Failed Undetermined Phase Cloud Retrieval Successful Liquid Water Cloud Retrieval Successful Ice Cloud Retrieval Successful Undetermined Phase Cloud Retrieval
1.6 - 2.1 μm PCL (Partly Cloudy) Cloud Retrieval Phase & Outcome <i>(The Cloud Retrieval Phase Flag and Outcome Flag can be read as a "combined" flag as documented here -- or read as separate flags -- the bit structure is identical. (See page 25 of this Plan for details))</i>	4	0 1 2 3 4 10 11 12	Cloud Mask Undetermined ^{n,f} Not Processed (typically clear) ^f Failed Liquid Water Cloud Retrieval Failed Ice Cloud Retrieval Failed Undetermined Phase Cloud Retrieval Successful Liquid Water Cloud Retrieval Successful Ice Cloud Retrieval Successful Undetermined Phase Cloud Retrieval

VNSWIR - 2.1 μm (Primary)	4	0	Cloud Mask Undetermined ^{n,f}
PCL (Partly Cloudy)		1	Not Processed (typically clear) ^f
Cloud Retrieval Phase & Outcome		2	Failed Liquid Water Cloud Retrieval
<i>(The Cloud Retrieval Phase Flag and</i>		3	Failed Ice Cloud Retrieval
<i>Outcome Flag can be read as a "combined"</i>		4	Failed Undetermined Phase Cloud Retrieval
<i>flag as documented here -- or read as separate</i>		10	Successful Liquid Water Cloud Retrieval
<i>flags -- the bit structure is identical. (See page</i>		11	Successful Ice Cloud Retrieval
<i>25 of this Plan for details)</i>		12	Successful Undetermined Phase Cloud Retrieval

ⁿ Cloud Optical Property retrieval not attempted

^f fill values used for Cloud Optical Property retrieval

Alternate Way to Read & Interpret *Cloud Retrieval Phase Flag* & *Cloud Retrieval Outcome Flag* “Pairs”

Some Cloud Optical Property QA flags always appear in adjacent pairs. One of the more important flag pairings is the **Cloud Retrieval Phase Flag** and the **Cloud Retrieval Outcome Flag**, which appear in a number of places in the 06_L2 SDS *Quality_Assurance_1km*. It should be noted that each of these flag pairs (example shown in the first table below) can be read as a single “combined” flag and interpreted as shown in the second table below. This combining is how the Level-3 code correctly computes the (*Successful*) *Cloud_Retrival_Fraction* and the (*Successful Retrieval*) *Pixel_Counts* statistics.

Further it should be noted that ALL of the **Cloud Retrieval Phase Flags** in the *Quality_Assurance_1km* SDS, contain identical information. That is, the Cloud Phase does NOT vary between the various Cloud Optical Property Retrieval Algorithms; however the adjacent **Retrieval Outcome Flags** contain information that varies depending on the Retrieval Algorithm. Therefore when you read these two flags - - one repeated (phase) and one unique (outcome) -- as a single “combined” flag, one gets unique information on successful and failed retrievals for each of the three cloud phases in each of the various cloud optical property retrieval algorithms. See the 2nd table below.

Finally note that the 7th through 9th (last 3) bytes of the *Quality_Assurance_1km* SDS (documented on the previous pages of this QA Plan) are shown as having the “combined flag” format displayed in the second table below. Note that you can devolve those combined flags into separate flags as described in the first table below. **These two formats are completely identical (and interchangeable) in terms of the bit structure of the QA array.**

Note that “Phase Flag” and “Outcome Flag” pairs in <i>Quality_Assurance_1km</i> ...			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Retrieval Phase Flag	3	0	Cloud Mask Undetermined
		1	Not Processed (typically clear)
		2	Liquid Water Cloud
		3	Ice Cloud
		4	Undetermined Phase Cloud
Cloud Retrieval Outcome Flag	1	0	Retrieval not attempted or unsuccessful
		1	Retrieval successful

... can be read and interpreted this way			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Retrieval Phase & Outcome Flag	4	0	Cloud Mask Undetermined
		1	Not Processed (typically clear)
		2	Failed Liquid Water Cloud Retrieval
		3	Failed Ice Cloud Retrieval
		4	Failed Undetermined Phase Cloud Retrieval
		10	Successful Liquid Water Cloud Retrieval
		11	Successful Ice Cloud Retrieval
		12	Successful Undetermined Phase Cloud Retrieval

Cloud Top Properties

Cloud Top Property QA flags are stored in 2 separate QA arrays (SDS's). The first SDS (*Cloud_Mask_5km*, which is 2 bytes in length) contains cloud mask QA flags, which are subsampled (every 5th line, every 5th pixel) from the Cloud Mask (35_L2) product.

It should be noted that the *Cloud_Mask_5km* SDS was expanded from 1 byte to 2 bytes for Collection 006. The first byte remains the same as in Collection 005; the second byte is a reworked combination of flags in byte one, which were needed to correctly compute a number of new Surface Type parameters in Level-3.

The second SDS (*Quality_Assurance_5km*, which is 10 bytes in length) contains the product quality flags, retrieval processing flags, and input data resource flags. Detail on the content of these two arrays is provided below. All Cloud Top Property QA Flag arrays have the following characteristics:

- Spatial resolution: 5 ° 5 km
- Processing mode: Both Day and Night

Scientific Data Set (SDS): "Cloud_Mask_5km" Description: Cloud mask 1 km QA flags subsampled at 5 km (every 5 th line, every 5 th pixel) Length: C006 = 2 bytes (16 bits) C005/051 = 1 bytes (8 bits)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Mask Status Flag	1	0 1	Undetermined Determined
Cloud Mask Cloudiness Flag	2	0 1 2 3	Confident Cloudy (or Fill, if Status Flag = 0) Probably Cloudy Probably Clear Confident Clear
Day / Night Flag	1	0 1	Night (or Fill, if Status Flag = 0) Day
Sunglint Flag	1	0 1	Yes (or Fill, if Status Flag = 0) No
Snow / Ice Flag	1	0 1	Yes (or Fill, if Status Flag = 0) No
Surface Type Flag	2	0 1 2 3	Ocean or Deep Lakes and Rivers (or Fill) Coast or Shallow Lakes and Rivers Desert Land

C6 Sunlint Flag (<i>Retooled</i>) (Note: "Sunlint = Yes" goes from 0 above to 2 here)	2	0	Fill or CTP retrieval fail
		1	No Sunlint & CTP retrieval success
		2	Sunlint & CTP retrieval success
C6 Snow / Ice Flag (<i>Retooled</i>) (Note: "Snow/Ice = Yes" goes from 0 above to 2 here)	2	0	Fill or CTP retrieval fail
		1	No Snow/Ice & CTP retrieval success
		2	Snow/Ice & CTP retrieval success ^f
C6 Surface Type Flag (<i>Retooled</i>) (Note 1: Surface Type Flags are re-indexed here, as compared to the above (original) S.T.Flag) Note 2: Cloud Mask Status Flag must be 1 for categories 1 through 5 because Maddux wants the surface type the cloud top properties were retrieved over	3	0	Fill or CTP retrieval fail
		1	Ocean, Deep Lakes and Rivers & CTP retr. success
		2	Coast, Shallow Lakes and Rivers & CTP retr. success
		3	Desert & CTP retrieval success
		4	Land & CTP retrieval success
		5	All other valid (non-fill) surface types & CTP retr. succ.
Day / Night Flag (Note this spare bit was changed to the Day/Night Flag in 2013. Flag needed to be repeated for L3 Surface Types.)	1	0	Night (or Fill, if Status Flag = 0)
		1	Day

^f **For Daytime:** Snow or ice under thin clouds only using spectral test, which cannot detect snow under thick clouds. Flag hardwired to snow over Antarctic land. **For Nighttime:** Snow or ice under all clouds (using ancillary snowmask data).

<i>Scientific Data Set (SDS): "Quality_Assurance_5km"</i> <i>Description: Cloud Top Property product quality and retrieval processing QA flags at 5x5 km</i> <i>Length: 10 bytes (80 bits)</i>			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Top Pressure Usefulness Flag	1	0	Not useful
		1	Useful
Cloud Top Pressure Confidence Flag	3	0	Fill
		1	Marginal Confidence (<i>not currently used</i>)
		2	Good Confidence (<i>not currently used</i>)
		3	Very Good Confidence
Cloud Top Temperature Usefulness Flag	1	0	Not useful
		1	Useful
Cloud Top Temperature Confidence Flag	3	0	Fill
		1	Marginal Confidence (<i>not currently used</i>)
		2	Good Confidence (<i>not currently used</i>)
		3	Very Good Confidence
Cloud Fraction Usefulness Flag	1	0	Not useful
		1	Useful
Cloud Fraction Confidence Flag	3	0	Fill
		1	Marginal Confidence (<i>not currently used</i>)
		2	Good Confidence (<i>not currently used</i>)
		3	Very Good Confidence
Cloud Effective Emissivity Usefulness Flag	1	0	Not useful
		1	Useful

Cloud Effective Emissivity Confidence Flag	3	0 1 2 3	Fill Marginal Confidence (<i>not currently used</i>) Good Confidence (<i>not currently used</i>) Very Good Confidence
Cloud Phase Infrared Usefulness Flag	1	0 1	Not useful Useful
Cloud Phase Infrared Confidence Flag	3	0 1 2 3	Fill Marginal Confidence (<i>mixed or undetermined phase</i>) Good Confidence (<i>not currently used</i>) Very Good Confidence (<i>ice or liquid water phase</i>)
<i>processing path flags</i>			
"Cirrus" Flag (CTP ≤ 680 hPa & CEE ≤ 0.95) <i>Note: Changed CTP threshold to 680 (from 700) for C006</i>	2	0 1 2 3	Missing Cloudy, No cirrus found Cloudy, Cirrus found Clear Sky (<i>New Category defined for C006</i>)
"High Cloud" Flag (CTP < 440 hPa) <i>Note: Changed CTP threshold to 440 (from 400) for C006</i>	2	0 1 2 3	Missing Cloudy, No high cloud found Cloudy, High cloud found Clear Sky (<i>New Category defined for C006</i>)
Number of Cloudy Pixels	Int 8	0-25	Number of 1 km cloudy pixels in 5x5 km retrieval area
Number of Clear Pixels	Int 8	0-25	Number of 1 km clear pixels in 5x5 km retrieval area
Number of Missing Pixels	Int 8	0-25	Number of 1 km missing pixels in 5x5 km retrieval area
<i>Additional QA and input data resource flags</i>			
Cloud Top Height Usefulness Flag	1	0 1	Not useful Useful
Cloud Top Height Confidence Flag	3	0 1 2 3	Fill Marginal Confidence (<i>not currently used</i>) Good Confidence (<i>not currently used</i>) Very Good Confidence
Overshooting Thunderstorm Top Flag	2	0 1 2	Fill No Overshooting Thunderstorm Top Found Overshooting Thunderstorm Top Found
Clear Radiance Origin	2	0 1 2	MOD35* (MODIS Cloud Mask) Forward calculation from model (NCEP GDAS) Other
Moisture Profile	2	0 1 2 3	NCEP GDAS GMAO AIRS / AMSU Other


Temperature Profile	2	0 1 2 3	NCEP GDAS GMAO AIRS / AMSU Other
Dir Surface Temperature Over Land	2	0 1 2 3	NCEP GDAS GMAO MOD11 [*] (MODIS Land Surface Temperature) Other
Surface Temperature Over Ocean	2	0 1 2 3	Reynolds blended GMAO MOD28 [*] (MODIS Sea Surface Temperature) Other
Surface Pressure	2	0 1 2	NCEP GDAS GMAO Other
Topography	2	0 1	EOS DEM Other
Surface Emissivity	2	0 1	CERES MOD11 [*] (MODIS Land Surface Temperature)
Surface Type	2	0 1 2 3	Loveland 1 km NA Olson Ecosystem MOD12 [*] (MODIS Land Cover) Other
<i>additional QA flags</i>			
Cloud Height Category	3	0 1 2 3 4 5	Fill (Missing data) Clear Sky or too clear for retrieval (CF = 0, 4, 8, 12%) ¹ Cloudy (CF ≥ 16%), CTP retrieval started, but failed ² Low Clouds (CTP ≥ 680 hPa) Middle Clouds (680 hPa > CTP ≥ 440 hPa) High Clouds (CTP < 440 hPa)
Nadir View Angle Flag	2	0 1 2	Fill (Missing data) Near Nadir View Angle (View Angle ≤ 32°) Oblique View Angle (View Angle > 32°)
Cloud Height Method (Note: Index's 0-4 and 6 were set to match the Cloud_Height_Method SDS. Index 5 is not used in the Cloud_Height_Method SDS, a new category 5 was added here)	3	0 1 2 3 4 5 6 7	Fill (Missing data) CO ₂ Slicing (band 36/35) CO ₂ Slicing (band 35/34) CO ₂ Slicing (band 35/33) CO ₂ Slicing (band 34/33) Cloudy, but not retrieved (retrieval failed) ² IR Window Clear Sky ¹

* MOD prefix denotes MODIS/Terra (AM overpass) data -- MYD prefix should be substituted for MODIS/Aqua (PM overpass) data.

¹ Clear means too clear for a retrieval. Only 0, 1, 2, or 3 pixels out of 25 tagged cloudy by CM (i.e. CF=0, 4, 8, 12%). Note "True cloud fraction" retained (clouds).

² Cloudy, but not retrieved (retrieval failed) in 3x3 grids where at least 4 pixels tagged Cloudy (i.e. CF ≥ 16%). This failure information was requested by Bruce Madden.

Sample Runtime QA Flag Images from the Cloud Product



¹ Note: "Thin Cirrus Flag" from Cloud Mask is suspect (due to 1.38 μ m test limitations) for portions of this granule.

Note: Images were created by "bitflag_visualizer", available at http://modis-atmos.gsfc.nasa.gov/tools_bitflag_visualizer.html.

Atmospheric Profile Product: MOD07_L2 (Terra) & MYD07_L2 (Aqua)

Atmospheric Profile QA flags are stored in 3 separate QA arrays (SDS's). The first SDS (*Cloud_Mask*, 1 byte in length) contains cloud mask QA flags, which are subsampled (every 5th line, every 5th pixel) from the Cloud Mask (35_L2) product. The second SDS (*Quality_Assurance*, 10 bytes in length) contains the product quality flags, retrieval processing flags, and and input data resource flags for all products except Water Vapor. The third SDS (*Quality_Assurance_Infrared*, 5 bytes in length) contains the product quality flags, retrieval processing flags, and and input data resource flags for Water Vapor (using the IR method). Detail on the content of these three arrays is provided below. All Atmospheric Profile QA Flag arrays have the following characteristics:

- Spatial resolution: 5 × 5 km
- Processing mode: Both Day and Night

Scientific Data Set (SDS): “Cloud_Mask” Description: Cloud Mask QA flags subsampled at 5 km (every 5th line, every 5th pixel) Length: 1 byte (8 bits)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Mask Status Flag	1	0	Undetermined
		1	Determined
Cloud Mask Cloudiness Flag	2	0	Confident Cloudy (or Fill, if Status Flag = 0)
		1	Probably Cloudy
		2	Probably Clear
		3	Confident Clear
Day / Night flag	1	0	Night (or Fill, if Status Flag = 0)
		1	Day
Sunglint Flag	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Snow / Ice Flag	1	0	Yes (or Fill, if Status Flag = 0)
		1	No
Surface Type Flag	2	0	Ocean or Deep Lakes and Rivers (or Fill)
		1	Coast or Shallow Lakes and Rivers
		2	Desert
		3	Land

Scientific Data Set (SDS): "Quality_Assurance"

Description: Atmospheric Profile product quality and retrieval processing QA flags at 5x5 km

Length: 10 bytes (80 bits)

Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Retrieved Temperature Profile Usefulness Flag ⁿ	1	0 1	Not useful Useful
Retrieved Temperature Profile Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality not currently used not currently used
Retrieved Moisture Profile Usefulness Flag ⁿ	1	0 1	Not useful Useful
Retrieved Moisture Profile Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality not currently used not currently used
Total Ozone Burden Usefulness Flag ⁿ	1	0 1	Not useful Useful
Total Ozone Burden Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality not currently used not currently used
Stability Indices (Lifted Index) Usefulness Flag ⁿ	1	0 1	Not useful Useful
Stability Indices (Lifted Index) Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality not currently used not currently used
Stability Indices (K Index) Usefulness Flag ⁿ	1	0 1	Not useful Useful
Stability Indices (K Index) Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality not currently used not currently used
Stability Indices (Total Totals) Usefulness Flag ⁿ	1	0 1	Not useful Useful
Stability Indices (Total Totals) Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality not currently used not currently used
<i>processing path flags</i>			
Number of Cloudy Pixels	Int 8	0-25	Number of 1km cloudy pixels in 5x5 km retrieval area

Number of Clear Pixels	Int 8	0-25	Number of 1km clear pixels in 5x5 km retrieval area
Number of Missing Pixels	Int 8	0-25	Number of 1km missing pixels in 5x5 km retrieval area
Method of Profiles Retrieval	2	0 1 2 3	Statistical Physical Other No retrieval
Method of Ozone Retrieval	2	0 1 2 3	RTE Perturbation Upper and Lower stratospheric ozone method Other No retrieval
Spares	4		TBD
<i>input data resource flags</i>			
Guess Moisture profile	2	0 1 2 3	NCEP GDAS GMAO AIRS / AMSU Other
Guess Temperature Profile	2	0 1 2 3	NCEP GDAS GMAO AIRS / AMSU Other
Surface Temperature Over Land	2	0 1 2	NCEP GDAS GMAO MOD11 * (MODIS Land Surface Temperature)
Surface Temperature Over Ocean	2	0 1 2 3	Reynolds blended GMAO MOD28 * (MODIS Sea Surface Temperature) Other
Surface Pressure	2	0 1 2	NCEP GDAS GMAO Other
Ozone First Guess Profile	2	0 1 2 3	TOMS TOVS GMAO Other
Spares	4		TBD
Spares	8		TBD


* MOD prefix denotes MODIS/Terra (AM overpass) data -- MYD prefix should be substituted for MODIS/Aqua (PM overpass) data.

ⁿ Flag is not reliable in Not Useful / Bad regions (i.e. Not Useful pixels are incorrectly tagged as Useful, etc. about 1/2 the time.)

Scientific Data Set (SDS): “Quality_Assurance_Infrared” Description: Product quality and retrieval processing flags for Precipitable Water (IR) at 5x5 km Length: 5 bytes (40 bits)			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Total Precipitable Water (IR) Usefulness Flag ⁿ	1	0 1	Not useful Useful
Total Precipitable Water (IR) Confidence Flag ⁿ	3	0 1 2 3	Fill (Bad or Cloudy) Best Quality <i>not currently used</i> <i>not currently used</i>
Spares	4		TBD
Number of Cloudy Pixels	Int 8	0-25	Number of 1 km Cloudy pixels n 5x5 retrieval area
Number of Clear Pixels	Int 8	0-25	Number of 1 km Clear pixels in 5x5 retrieval area
Number of Missing Pixels	Int 8	0-25	Number of 1 km Missing pixels in 5x5 retrieval area
IR Water Vapor Retrieval Method Used	2	0 1 2 3	Split Window (11-12 μm) Technique Integration of Moisture Profile Other No Retrieval
Spares	6		TBD

ⁿ Flag is not reliable in Not Useful / Bad regions (i.e. Not Useful pixels are incorrectly tagged as Useful, etc. about ½ the time.)

Sample Runtime QA Flag Images from the Atmospheric Profile Product



¹ All 07_L2 Atmospheric Profile Usefulness (and Confidence) Flags are noisy.

Collection 006 Suggestion: Remove noise from all Usefulness and Confidence Flags in the 07_L2 Atmospheric Profile product.

Note: Images were created by "bitflag_visualizer", available at http://modis-atmos.gsfc.nasa.gov/tools_bitflag_visualizer.html.

Note: The local version of the Cloud Mask QA flags in the 07_L2 Atmospheric Profile product are stored in an SDS named "Cloud_Mask." Users should note that this array is different from the primary "Cloud_Mask" array stored in 35_L2 Cloud Mask product, which has the same name.

Cloud Mask Product: MOD35_L2 (Terra) & MYD35_L2 (Aqua)

For the Cloud Mask product, the QA flags are stored in a single ten byte SDS (called *Quality_Assurance*) containing the product quality flags, retrieval processing flags and input data resource flags. The Cloud Mask QA Flag array has the following characteristics:

- Spatial resolution: 1×1 km and 250×250 m
- Processing mode: Both Day and Night

Scientific Data Set (SDS): <i>"Quality_Assurance"</i> Description: <i>Cloud Mask QA flags at 1 km and 250 m</i> Length: <i>10 bytes (80 bits)</i>			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Mask (1 km) Usefulness Flag	1	0 1	Not useful Useful
Cloud Mask (1 km) Confidence Flag	3	0 1 2 3 4 5 6 7	Lowest Confidence Not Used Not Used Not Used Intermediate Confidence Not Used High Confidence Highest Confidence
Spares	4		TBD
<i>individual test application flags</i>			
Non Cloud Obstruction (NCO) Test <i>(Spectral tests for fire, smoke, & dust)</i>	1	0 1	Not Applied Applied
Thin Cirrus (Solar) Test	1	0 1	Not Applied Applied
Snow Cover from Ancillary Data	1	0 1	Ancillary Data Not Tested Ancillary Data Tested
Thin Cirrus (IR) Test	1	0 1	Not Applied Applied
Cloud Adjacency Test <i>(nearest neighbor pixels)</i>	1	0 1	Not Applied Applied
IR Threshold Test	1	0 1	Not Applied Applied
High Cloud (CO ₂) Test	1	0 1	Not Applied Applied
High Cloud (6.7 μm) Test	1	0 1	Not Applied Applied

High Cloud (1.38 μm) Test	1	0 1	Not Applied Applied
High Cloud (3.9-12 μm) Test	1	0 1	Not Applied Applied
Transmissive High Cloud (11-12 μm) Test	1	0 1	Not Applied Applied
3.9-11 μm Test	1	0 1	Not Applied Applied
0.412/0.68/0.86 μm Reflectance Test	1	0 1	Not Applied Applied
0.86/0.68 Ratio Test	1	0 1	Not Applied Applied
Clear Sky Restoral Test (Coastal NDVI)	1	0 1	Not Applied Applied
7.3-11 μm Test (night land, night snow, polar night)	1	0 1	Not Applied Applied
Ocean 8.6-11 μm Test	1	0 1	Not Applied Applied
Clear Sky Restoral Test (Spatial variability, water)	1	0 1	Not Applied Applied
Clear Sky Restoral Tests (Polar night, land, sun-glint)	1	0 1	Not Applied Applied
Surface Temperature Test	1	0 1	Not Applied Applied
Suspended Dust Test	1	0 1	Not Applied Applied
Night Ocean 8.6-7.3 μm Test	1	0 1	Not Applied Applied
Night Ocean 11 μm Spatial Variability Test	1	0 1	Not Applied Applied
Night Ocean Low Cloud Test (3.9-11 μm)	1	0 1	Not Applied Applied
250 m Visible Tests (Repeated 16 times)	1 (16)	0 1	Not Applied Applied
<i>input data information flags</i>			
Number of bands used to generate cloud mask	2	0 1 2 3	None 1-7 8-14 15-21
Number of spectral tests used to generate cloud mask	2	0 1 2 3	None 1-3 4-6 7-9
Spares	4		TBD

<i>input data resource flags</i>			
Clear Radiance Origin	2	0 1 2	MOD35* (MODIS Cloud Mask) Forward calculation from NCEP GDAS model Other
Surface Temperature Over Land	2	0 1 2 3	NCEP GDAS GMAO MOD11* (MODIS Land Surface Temperature) Other
Surface Temperature Over Ocean	2	0 1 2 3	Reynolds blended GMAO MOD28* (MODIS Sea Surface Temperature) Other
Surface Winds	2	0 1 2	NCEP GDAS GMAO Other
Ecosystem Map	2	0 1 2 3	Loveland NA 1 km Olson Ecosystem MOD12* (MODIS Land Cover) Other
Snow Mask	2	0 1 2	MOD33* (MODIS Snow Cover) SSM/I product Other
Ice Cover	2	0 1 2	MOD42* (MODIS Sea Ice) SSM/I product Other
Land/Sea Mask	2	0 1 2	USGS 1 km 6 level USGS 1 km binary Other
Digital Elevation Model	1	0 1	EOS DEM Not used
Precipitable Water	2	0 1 2	NCEP GDAS GMAO MOD07* (MODIS Atmospheric Profile)
Spares	5		TBD

* MOD prefix denotes MODIS/Terra (AM overpass) data -- MYD prefix should be substituted for MODIS/Aqua (PM overpass) data.


Note: All Cloud Mask Tests are described in the Cloud Mask Algorithm Theoretical Basis Document (ATBD).

Sample Runtime QA Flag Images from the Cloud Mask Product

Reference L1B Image of Scene

12 Feb 2001 15:10

Flag = "Shadow Detection Test"



Note: Images were created by "bitflag_visualizer", available at http://modis-atmos.gsfc.nasa.gov/tools_bitflag_visualizer.html.

Joint Atmosphere Product: MODATML2 (Terra) & MYDATML2 (Aqua)

The *post-launch* Level 2 (L2) Joint Atmosphere product contains a spectrum of key parameters gleaned from the complete set of standard *at-launch* L2 products: Aerosol, Water Vapor, Cloud, Profile, and Cloud Mask. The L2 Joint Atmosphere product was designed to provide a convenient way to obtain the “greatest hits” of MODIS Atmosphere within a single HDF file – small enough to minimize data transfer and storage requirements, yet robust enough to be useful to a significant number of MODIS data users. Scientific data sets (SDS’s) contained within the L2 Joint Atmosphere product are stored at 5 km and 10 km (at nadir) spatial resolutions.

It should be noted that standard L2 parameters, which are retrieved at 1 km resolution, are subsampled at 5 km (every 5th line, every 5th pixel) for inclusion in the L2 Joint Atmosphere Product. This means that the *Cloud_Quality_Assurance* SDS (detailed below) matches exactly the *Quality_Assurance_1km* SDS contained within the Cloud (06_L2) product, except that it is sub-sampled at 5 km resolution. The *Aerosol_Quality_Assurance* array (also detailed below) is stored at its native resolution of 10 km, however it only contains a small subset of the flags found within the source *Quality_Assurance_Land* and *Quality_Assurance_Ocean* SDS’s in the Aerosol (04_L2) product. All Joint Atmosphere QA Flag arrays have the following characteristics:

- Spatial resolution: 5x5 km (Cloud Optical Properties) & 10x10 km (Aerosol)
- Processing mode: Daytime only

Scientific Data Set (SDS): “ <i>Cloud_Quality_Assurance</i> ” Description: <i>Cloud Optical Property QA flags subsampled at 5 km resolution</i> Length: <i>5 bytes (40 bits)</i>			
Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Cloud Optical Thickness (τ) Usefulness Flag	1	0 1	Not useful Useful
Cloud Optical Thickness (τ) Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
Cloud Optical Thickness (τ) Out-of-Bounds Flag	2	0 1 2	Within bounds ($0 < \tau \leq 100$) or Fill $100 < \tau \leq 150$ $\tau > 150$
Cloud Effective Radius Usefulness Flag	1	0 1	Not useful Useful
Cloud Effective Radius Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence

Cloud Water Path Usefulness Flag	1	0 1	Not useful Useful
Cloud Water Path Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
1.6 / 2.1 μm Cloud Retrieval Phase Flag <i>Note: Retrieval performed only over ocean, snow, & ice. Cloud phase determined by Cloud Mask, IR, & SWIR-based decision tree.</i>	3	0 1 2 3 4	Cloud Mask Undetermined or Non-Snow Land ^{n,f} Not Processed (typ. clear) over Ocean, Snow, Ice ^f Liquid Water Cloud Ice Cloud Undetermined Phase Cloud
1.6 / 2.1 μm Cloud Retrieval Outcome	1	0 1	1.6 / 2.1 μm retrieval not attempted or unsuccessful ^f 1.6 / 2.1 μm retrieval successful
Spare	1		TBD
Primary Cloud Retrieval Phase Flag <i>Note: Cloud phase determined by Cloud Mask, IR, & SWIR-based decision tree.</i>	3	0 1 2 3 4	Cloud Mask Undetermined (missing) ^{n,f} Not Processed (typically clear) ^f Liquid Water Cloud Ice Cloud Undetermined Phase Cloud
Primary Cloud Retrieval Outcome	1	0 1	Retrieval not attempted or unsuccessful ^f Retrieval successful
Rayleigh Correction	1	0 1	No Yes
Atmospheric Correction	1	0 1	No Yes
Band Used for Primary Optical Thickness Retrieval	2	0 1 2 3	Retrieval not attempted ^f 0.645 μm (land) 0.858 μm (water) 1.24 μm (snow / ice)
1.6 / 2.1 μm Cloud Optical Thickness (τ) Usefulness Flag	1	0 1	Not useful Useful
1.6 / 2.1 μm Cloud Optical Thickness (τ) Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
1.6 / 2.1 μm Cloud Effective Radius Usefulness Flag	1	0 1	Not useful Useful
1.6 / 2.1 μm Cloud Effective Radius Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence

Clear Sky Restoral	2	0 1 2 3	Not Restored Restored via Edge Detection Restored via Spatial Variance Restored via 250 m Tests
1.6 / 2.1 μm Cloud Water Path Usefulness Flag	1	0 1	Not useful Useful
1.6 / 2.1 μm Cloud Water Path Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
Primary Cloud Retrieval Multilayer Cloud & Phase Flag	3	0 1 2 3 4 5 6 7	Cloud mask undetermined (missing) ^{n,f} Not Processed (typically clear) ^f Single Layer Liquid Water Cloud Multi Layer Liquid Water Cloud Single Layer Ice Cloud Multi Layer Ice Cloud Single Layer Undetermined Phase Cloud Multi Layer Undetermined Phase Cloud
Primary Cloud Retrieval Outcome (flag duplicated from 3 rd byte)	1	0 1	Retrieval not attempted or unsuccessful ^f Retrieval successful
Spare	1		TBD

ⁿ Cloud Optical Property retrieval not attempted

^f fill values used for Cloud Optical Property retrieval

Scientific Data Set (SDS): "Aerosol_Quality_Assurance"

Description: Aerosol QA flags (a small subset of flags only) at 10x10 km resolution

Length: 1 byte (8 bits)

Flag Name	Number of Bits	Bit Values	Bit Value Definitions
Aerosol Optical Depth Land 0.47 μm Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
Aerosol Optical Depth Land 0.66 μm Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
Aerosol Optical Depth Ocean Avg. Solution Confidence Flag	2	0 1 2 3	No Confidence or Fill Marginal Confidence Good Confidence Very Good Confidence
Spares	2		TBD

Appendix B:

Inventory Metadata

of

Atmosphere Products

Introduction

This appendix describes the Inventory Metadata for all MODIS Atmosphere products. Inventory Metadata are searchable strings, stored within each HDF file, that can be used to assist users to select granules that meet predetermined criteria. Inventory Metadata are stored in the HDF global attribute "CoreMetadata.0".

There are two main types of Inventory Metadata:

- ***ECS (or Core) Inventory Metadata*** that can be found, in the same format, in all MODIS Atmosphere products. This includes strings that specify *Date/Time Range* and *Bounding Rectangle* (geolocation). Note that ECS stands for EOSDIS Core System.
- ***QA Inventory Metadata***, which is product-specific (i.e.: varies from product to product), and typically provides more scientifically revealing information about a granule.

The primary data archive and ordering system that utilizes the full set of both ECS and QA Inventory Metadata to search and sort MODIS HDF granules is the ***EOS Data Gateway (EDG)***, (<http://redhook.gsfc.nasa.gov/ims-bin/pub/nph-ims.cgi>).

Other data archive and ordering systems utilize only a small subset of the ECS (Core) Inventory Metadata. This subset of Core Inventory Metadata utilized typically includes only the *Date/Time Range* and *Bounding Rectangle* (geolocation) strings. These less powerful, but perhaps more user-friendly, data archive and ordering systems include the ***MODIS Multiple Data Ordering Page (MDOP)***, (http://disc.gsfc.nasa.gov/daac-bin/MODIS/Data_order.pl?PRINT=1) and the ***Atmosphere Archive and Distribution System - Web interface (AADSWeb)***, (<http://aadsweb.nascom.nasa.gov>).

ECS Inventory Metadata

ECS (or Core) Inventory Metadata is high-level information that can be used to help sort MODIS HDF data files based on basic user requirements. Shown in the table below is the Core Inventory Metadata implemented in all (both Level 2 and Level 3) MODIS Atmosphere products. It includes information on granule location, collection day and time, orbit information, percent missing data, science quality and automatic (operational) quality flags specifically defined to be associated with the measured parameters of a product.

The automatic quality flag is set by the MODIS processing facility (either the Goddard DAAC (Distributed Active Archive Center) or MODAPS) at the time the granule is produced. The science quality flag is assigned initially by the science team, and then updated through an update utility, if desired. Both of these flags are associated with a particular retrieval algorithm and apply to all derived SDSs.

All MODIS Atmosphere Products

ECS (or Core) Inventory Metadata

ECS Core Attribute Name	Sample Value (shown for a L2 Joint Product)
LOCALGRANULEID	"MODATML2.A2001043.1510.005.2005105195411.hdf"
PRODUCTIONDATETIME	"2005-04-15T19:54:11.000Z"
DAYNIGHTFLAG	"Day"
LOCALVERSIONID	"005"
REPROCESSINGACTUAL	"processed once"
REPROCESSINGPLANNED	"further update is anticipated"
SHORTNAME	"MODATML2"
VERSIONID	"5"
PGEVERSION	"5.1.0"
INPUTPOINTER †	"MOD03.A2001043.1510.005.2005105143551.hdf, MOD04_L2.A2001043.1510.005.2005105193145.hdf, MOD05_L2.A2001043.1510.005.2005105193042.hdf, MOD06_L2.A2001043.1510.005.2005105195333.hdf, MOD07_L2.A2001043.1510.005.2005105171354.hdf, MOD35_L2.A2001043.1510.005.2005105171245.hdf"
<i>Date / Time Range</i>	
RANGEBEGINNINGTIME	15:10:00.000000
RANGEENDINGTIME	15:15:00.000000
RANGEBEGINNINGDATE	2001-02-12
RANGEENDINGDATE	2001-02-12
<i>Bounding Rectangle</i>	
EASTBOUNDINGCOORDINATE	-58.4177095990599
WESTBOUNDINGCOORDINATE	-85.5363614515652
NORTHBOUNDINGCOORDINATE	-11.9871990497647
SOUTHBOUNDINGCOORDINATE	-33.0740234458278
<i>Orbit Calculated Spatial Domain</i>	
ORBITNUMBER.1	6149
EQUATORCROSSINGLONGITUDE.1	-66.1949490630604
EQUATORCROSSINGDATE.1	2001-02-12
EQUATORCROSSINGTIME.1	15:06:12.273251
<i>Measured Parameter</i>	
PARAMETERNAME.1	"Cloud_Top_Pressure" (example)
SCIENCEQUALITYFLAG.1	"Not Investigated"
SCIENCEQUALITYFLAGEXPLANATION.1	"See http://modis-atmos.gsfc.nasa.gov/validation.html for more details on MODIS Atmosphere data quality."
AUTOMATICQUALITYFLAG.1	"Passed"
AUTOMATICQUALITYFLAGEXPLANATION.1	"Passed: >10% useable; Failed: <10% useable"
QAPERCENTMISSINGDATA.1 #	23

† MODIS product input files # Pertains to Measured ParameterName and refers to 23% of granule being clear (no CTP retrieval)

QA Inventory Metadata

In addition to the ECS (or Core) Inventory Metadata described above, product-specific **QA Inventory Metadata** is used to accommodate the QA needs of each specific Level 2 (only) MODIS Atmosphere product. There is no QA Inventory Metadata defined or stored for Level 3 Atmosphere products.

The QA Inventory Metadata includes statistics reported by either the MODIS cloud mask algorithm or by the product generation algorithm itself. For all L2 MODIS atmosphere products, the QA Inventory Metadata may include:

- % successful rate of retrieval
- % low confidence clear sky
- % various cloud types
- % non-cloud obstruction
- % day processed
- % night processed
- maximum solar zenith angle
- minimum solar zenith angle
- % land processed
- % water processed
- % snow background processed
- % shadow processed
- % sunglint processed

An important feature of QA Inventory Metadata is its searchability. A user can skip an entire granule of data if the requested criteria are not satisfied, or can examine the granule in more detail if criteria are satisfied. Detailed descriptions of product-specific QA Inventory Metadata for each MODIS Atmosphere product follows.

Aerosol Product: MOD04_L2 (Terra) & MYD04_L2 (Aqua)

QA Inventory Metadata

(Note: Denominator in all Percentages = Total Pixels in entire Granule)

Field Name	Field Description	Sample Value [^]
<i>Aerosol Retrieval</i>		
AdditionalAttributeName.1	SuccessfulRetrievalPct_Land ¹	9.10
AdditionalAttributeName.2	SuccessfulRetrievalPct_Ocean ²	15.77
<i>Cloud and Non Cloud Obstructions</i>		
AdditionalAttributeName.3	LowConfidentClearPct ^m	69.15
AdditionalAttributeName.11	ThinCirrusSolarFoundPct ^m	22.56
AdditionalAttributeName.12	ThinCirrusIR_FoundPct ^m	3.48
AdditionalAttributeName.13	NonCloudObstructionFoundPct ^m	0.25
<i>Day/Night</i>		
AdditionalAttributeName.4	DayProcessedPct ^m	100.00
AdditionalAttributeName.5	NightProcessedPct ^m	0.00
<i>Solar Zenith Angle</i>		
AdditionalAttributeName.14	MaxSolarZenithAngle ^m	40.53
AdditionalAttributeName.15	MinSolarZenithAngle ^m	14.10
<i>Scene Surface Type</i>		
AdditionalAttributeName.8	LandProcessedPct ^m	51.99
AdditionalAttributeName.9	WaterProcessedPct ^m	48.01
AdditionalAttributeName.7	Snow_IceSurfaceProcessPct ^m	0.00
AdditionalAttributeName.10	ShadowFoundPct ^m	0.08
AdditionalAttributeName.6	SunlintProcessPct ^{m, i}	48.07


¹ For the Aerosol over Land algorithm, only clear-sky, dark-surface land pixels are retrieved.

² For the Aerosol over Ocean algorithm, only clear-sky, non-glnt ocean pixels are retrieved.

^m Fields are COPIED from the Cloud Mask QA Inventory Metadata and apply to a 1x1 km resolution analysis of granule.

ⁱ Incorrectly computed (based on solar and viewing geometry only -- did not consider whether the pixel is over land or water).

[^] Sample Values from MODIS/Terra 12 Feb 2001 15:10



Water Vapor Product: MOD05_L2 (Terra) & MYD05_L2 (Aqua)

QA Inventory Metadata

(Note: Denominator in all Percentages = Total Pixels in entire Granule)

Field Name	Field Description	Sample Value [^]
<i>Water Vapor Retrieval</i>		
AdditionalAttributeName.1	SuccessfulRetrievalPct_NIR ¹	99.62
AdditionalAttributeName.2	SuccessfulRetrievalPct_IR ²	30.75
<i>Cloud and Non Cloud Obstructions</i>		
AdditionalAttributeName.3	LowConfidentClearPct ^m	69.15
AdditionalAttributeName.11	ThinCirrusSolarFoundPct ^m	22.56
AdditionalAttributeName.12	ThinCirrusIR_FoundPct ^m	3.48
AdditionalAttributeName.13	NonCloudObstructionFoundPct ^m	0.25
<i>Day/Night</i>		
AdditionalAttributeName.4	DayProcessedPct ^m	100.00
AdditionalAttributeName.5	NightProcessedPct ^m	0.00
<i>Solar Zenith Angle</i>		
AdditionalAttributeName.14	MaxSolarZenithAngle ^m	40.53
AdditionalAttributeName.15	MinSolarZenithAngle ^m	14.10
<i>Scene Surface Type</i>		
AdditionalAttributeName.8	LandProcessedPct ^m	51.99
AdditionalAttributeName.9	WaterProcessedPct ^m	48.01
AdditionalAttributeName.7	Snow_IceSurfaceProcessPct ^m	0.00
AdditionalAttributeName.10	ShadowFoundPct ^m	0.08
AdditionalAttributeName.6	SunglintProcessPct ^{m, i}	48.07

¹ For the Near IR Water Vapor algorithm, all non-missing pixels are retrieved (but only bright-land & ocean-glint pixels are reliable).


Collection 006 Suggestion: Fix SuccessfulRetrievalPct_NIR computation so the numerator = "reliable (useful) retrievals only".

² For the IR Water Vapor algorithm, only clear-sky pixels (over both land and ocean) are retrieved.

^m Fields are COPIED from the Cloud Mask QA Inventory Metadata and apply to a 1x1 km resolution analysis of granule.

ⁱ Incorrectly computed (based on solar and viewing geometry only -- did not consider whether the pixel is over land or water).

[^] Sample Values from MODIS/Terra 12 Feb 2001 15:10



Cloud Product: MOD06_L2 (Terra) & MYD06_L2 (Aqua)

QA Inventory Metadata


(Note: Denominator in all Percentages = Total Pixels in entire Granule)

Field Name	Field Description	Sample Value ^
<i>Cloud Top Property Retrieval</i>		
AdditionalAttributeName.1	SuccessCloudTopPropRtrPct_IR ¹	77.12
AdditionalAttributeName.2	SuccessCloudPhaseRtrPct_IR ¹	76.75
AdditionalAttributeName.4	LowCloudDetectedPct_IR	19.77
AdditionalAttributeName.5	MidCloudDetectedPct_IR	38.91
AdditionalAttributeName.6	HighCloudDetectedPct_IR	18.44
AdditionalAttributeName.8	ThickCloudDetectedPct_IR	28.21
AdditionalAttributeName.7	ThinCloudDetectedPct_IR	6.44
AdditionalAttributeName.9	OpaqueCloudDetectedPct_IR	42.47
AdditionalAttributeName.10	CirrusCloudDetectedPct_IR	34.65
AdditionalAttributeName.12	WaterCloudDetectedPct_IR	49.69
AdditionalAttributeName.11	IceCloudDetectedPct_IR	13.05
AdditionalAttributeName.13	MixedCloudDetectedPct_IR	4.86
AdditionalAttributeName.14	CloudPhaseUncertainPct_IR	9.15
<i>Cloud Optical Property Retrieval</i>		
AdditionalAttributeName.3	SuccessCloudOptPropRtrPct_VIS (<i>successful retrievals only</i>)	64.32
AdditionalAttributeName.18	CloudCoverFractionPct_VIS (<i>successful & unsuccessful</i>)	67.56
AdditionalAttributeName.19	WaterCloudDetectedPct_VIS (<i>successful & unsuccessful</i>)	45.54
AdditionalAttributeName.20	IceCloudDetectedPct_VIS (<i>successful & unsuccessful</i>)	20.86
<i>Scene Surface Type</i>		
AdditionalAttributeName.16	LandCoverFractionPct ^m	51.99
AdditionalAttributeName.15	OceanCoverFractionPct ^m	48.01
AdditionalAttributeName.17	SnowCoverFractionPct ^m	0.00

¹ Only 5x5 km grids determined to be cloudy are retrieved. CTP algorithms "detect" more clouds than Cloud Mask and Optical Prop. algorithms due to 5x5 avg'ing. CloudTopPropertyIR and CloudPhaseIR pcts. are different due to different bands being used.

^m Fields are COPIED from the Cloud Mask QA Inventory Metadata and apply to a 1x1 km resolution analysis of granule.

[^] Sample Values from MODIS/Terra 12 Feb 2001 15:10



Atmospheric Profile Product: MOD07_L2 (Terra) & MYD07_L2 (Aqua)

QA Inventory Metadata

(Note: Denominator in all Percentages = Total Pixels in entire Granule)


Field Name	Field Description	Sample Value [^]
<i>Atmospheric Profile Retrieval</i>		
AdditionalAttributeName.1	SuccessfulRetrievalPct ¹	30.75
<i>Cloud and Non Cloud Obstructions</i>		
AdditionalAttributeName.2	LowConfidenceClearPct ^m	69.15
AdditionalAttributeName.10	ThinCirrusSolarFoundPct ^m	22.56
AdditionalAttributeName.11	ThinCirrusIR_FoundPct ^m	3.48
AdditionalAttributeName.12	NonCloudObstructionFoundPct ^m	0.25
<i>Day/Night</i>		
AdditionalAttributeName.3	DayProcessedPct ^m	100.00
AdditionalAttributeName.4	NightProcessedPct ^m	0.00
<i>Solar Zenith Angle</i>		
AdditionalAttributeName.13	MaxSolarZenithAngle ^m	40.53
AdditionalAttributeName.14	MinSolarZenithAngle ^m	14.10
<i>Scene Surface Type</i>		
AdditionalAttributeName.7	LandProcessedPct ^m	51.99
AdditionalAttributeName.8	WaterProcessedPct ^m	48.01
AdditionalAttributeName.6	Snow_IceSurfaceProcessPct ^m	0.00
AdditionalAttributeName.9	ShadowFoundPct ^m	0.08
AdditionalAttributeName.5	SunglintProcessPct ^{m, i}	48.07

¹ For the Atmospheric Profiles algorithm, only clear-sky pixels (either land or ocean) are retrieved.

^m Fields are COPIED from the Cloud Mask QA Inventory Metadata and apply to a 1x1 km resolution analysis of granule.

ⁱ Incorrectly computed (based on solar and viewing geometry only -- did not consider whether the pixel is over land or water).

[^] Sample Values from MODIS/Terra 12 Feb 2001 15:10



Cloud Mask Product: MOD35_L2 (Terra) & MYD35_L2 (Aqua)

QA Inventory Metadata

(Note: Denominator in all Percentages = Total Pixels in entire Granule)


Field Name	Field Description	Sample Value [^]
<i>Cloud Mask Retrieval</i>		
AdditionalAttributeName.1	SuccessfulRetrievalPct ¹	100.00
<i>Cloud and Non Cloud Obstructions</i>		
AdditionalAttributeName.7	ClearPct250m	34.90
AdditionalAttributeName.6	CloudCoverPct250m	65.10
AdditionalAttributeName.2	VeryHighConfidenceClearPct	23.78
AdditionalAttributeName.3	HighConfidenceClearPct	5.09
AdditionalAttributeName.4	UncertainConfidentClearPct	1.98
AdditionalAttributeName.5	LowConfidentClearPct	69.15
AdditionalAttributeName.15	ThinCirrusSolarFoundPct	22.56
AdditionalAttributeName.16	ThinCirrusIR_FoundPct	3.48
AdditionalAttributeName.17	NonCloudObstructionFoundPct	0.25
<i>Day/Night</i>		
AdditionalAttributeName.8	DayProcessedPct	100.00
AdditionalAttributeName.9	NightProcessedPct	0.00
<i>Solar Zenith Angle</i>		
AdditionalAttributeName.18	MaxSolarZenithAngle	40.53
AdditionalAttributeName.19	MinSolarZenithAngle	14.10
<i>Scene Surface Type</i>		
AdditionalAttributeName.12	LandProcessedPct	51.99
AdditionalAttributeName.13	WaterProcessedPct	48.01
AdditionalAttributeName.11	Snow_IceSurfaceProcessPct	0.00
AdditionalAttributeName.14	ShadowFoundPct	0.08
AdditionalAttributeName.10	SunglintProcessPct ⁱ	48.07

¹ For the Cloud Mask algorithm, all non-missing L1B pixels are retrieved.

ⁱ Incorrectly computed (based on solar and viewing geometry only -- did not consider whether the pixel is over land or water).

Collection 006 Suggestion: Fix the SunglintProcessPct computation (noted above). Fix will propagate to other L2 products.

[^] Sample Values from MODIS/Terra 12 Feb 2001 15:10



Joint Atmosphere Product: MODATML2 (Terra) & MYDATML2 (Aqua)

QA Inventory Metadata

(Note: Denominator in all Percentages = Total Pixels in entire Granule)

Field Name	Field Description	Sample Value ^
<i>Aerosol Retrieval</i>		
AdditionalAttributeName.18	SuccessfulRetrievalPct_Land ¹	9.10
AdditionalAttributeName.19	SuccessfulRetrievalPct_Ocean ²	15.77
<i>Water Vapor Retrieval</i>		
AdditionalAttributeName.20	SuccessfulRetrievalPct_NIR ³	99.62
AdditionalAttributeName.21	SuccessfulRetrievalPct_IR ⁴	30.75
<i>Cloud Top Property Retrieval</i>		
AdditionalAttributeName.22	SuccessCloudTopPropRtrPct_IR ⁵	77.12
AdditionalAttributeName.23	SuccessCloudPhaseRtrPct_IR ⁵	76.75
<i>Cloud Optical Property Retrieval</i>		
AdditionalAttributeName.24	SuccessCloudOptPropRtrPct_VIS (<i>successful retrievals only</i>)	64.32
AdditionalAttributeName.25	CloudCoverFractionPct_VIS (<i>successful & unsuccessful</i>)	67.56
AdditionalAttributeName.26	WaterCloudDetectedPct_VIS (<i>successful & unsuccessful</i>)	45.54
AdditionalAttributeName.27	IceCloudDetectedPct_VIS (<i>successful & unsuccessful</i>)	20.86
<i>Cloud Mask Retrieval</i>		
AdditionalAttributeName.1	SuccessfulRetrievalPct ^m	100.00
<i>Cloud and Non Cloud Obstructions</i>		
AdditionalAttributeName.12	VeryHighConfidentClearPct ^m	23.78
AdditionalAttributeName.13	HighConfidentClearPct ^m	5.09
AdditionalAttributeName.14	UncertainConfidentClearPct ^m	1.98
AdditionalAttributeName.15	LowConfidentClearPct ^m	69.15
AdditionalAttributeName.16	ThinCirrusSolarFoundPct ^m	22.56
AdditionalAttributeName.17	ThinCirrusIR_FoundPct ^m	3.48
AdditionalAttributeName.9	NonCloudObstructionFoundPct ^m	0.25
<i>Day/Night</i>		
AdditionalAttributeName.2	DayProcessedPct ^m	100.00
AdditionalAttributeName.3	NightProcessedPct ^m	0.00
<i>Solar Zenith Angle</i>		
AdditionalAttributeName.10	MaxSolarZenithAngle ^m	40.53
AdditionalAttributeName.11	MinSolarZenithAngle ^m	14.10
<i>Scene Surface Type</i>		
AdditionalAttributeName.4	LandProcessedPct ^m	51.99
AdditionalAttributeName.5	WaterProcessedPct ^m	48.01
AdditionalAttributeName.6	Snow_IceSurfaceProcessedPct ^m	0.00
AdditionalAttributeName.7	ShadowFoundPct ^m	0.08
AdditionalAttributeName.8	SunlintProcessedPct ^{m, i}	48.07

¹ For Aerosol over Land algorithm, only clear-sky, dark-surface land pixels are retrieved.

² For Aerosol over Ocean algorithm, only clear-sky, non-glnt ocean pixels are retrieved.

³ For Near IR Water Vapor, all non-missing pixels are retrieved (but only bright-land & ocean-glnt pixels are reliable).


⁴ For IR Water Vapor, only clear-sky pixels (land and ocean) are retrieved. (Note: This number matches Atm Profile SuccessPct).

⁵ Only 5x5 km grids determined to be cloudy are retrieved. CTP algorithms “detect” more clouds than Cloud Mask and Optical Prop. algorithms due to 5x5 avg'ing. CloudTopPropertyIR and CloudPhaseIR pcts. are different due to different bands being used.

^m Fields are COPIED from the Cloud Mask QA Inventory Metadata and apply to a 1x1 km resolution analysis of granule.

ⁱ Incorrectly computed (based on solar and viewing geometry only -- did not consider whether the pixel is over land or water).

[^] Sample Values from MODIS/Terra 12 Feb 2001 15:10



Appendix C:

Archive Metadata

of

Atmosphere Products

Introduction

This appendix describes the Archive Metadata for all MODIS Atmosphere products. Archive Metadata, stored in the HDF global attribute "ArchiveMetadata.0", are designed to report summary statistics and information pertaining to a granule for documentation purposes only.

Archive metadata is stored (archived) along with science data sets of each granule. These values are not searchable. Archive Metadata is defined in both Level 2 and Level 3 MODIS Atmosphere products.

Arguably, the two most useful pieces of information in the Archive Metadata are the *LocalInputGranuleID* and *ProductionHistory* strings.

- The *LocalInputGranuleID* contains a full list of input files used to create the HDF product file in question. Input files can include other MODIS HDF product files (typically L1B Calibrated Radiances and/or L2 Cloud Mask) and additional ancillary files (such as algorithm specific look-up libraries, and/or snow, albedo and ecosystem maps) needed to run the L2 product algorithm.
- The *ProductionHistory* contains the program executable (PGE) version that was used to produce the HDF product file. This allows users to trace the maturity and quality of HDF data products. Specific changes incorporated into various versions of PGEs can be found in the Products section of the MODIS Atmosphere web site at http://modis-atmos.gsfc.nasa.gov/products_calendar_pge03.html (Note that the link specified is for PGE03, which produces the Cloud Mask and Atmospheric Profiles products. All other MODIS Atmosphere PGE's can be selected through the menu on the left hand side of this page).

The complete set of archive metadata for each MODIS Atmosphere product is described below.

Aerosol Product: MOD04_L2 (Terra) & MYD04_L2 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Aerosol 5-Min L2 Swath 10km"
PROCESSINGENVIRONMENT	"Linux minion5027 2.4.20-37.7.legacysmp #1 SMP Mon Sep 27 21:38:15 EDT 2004 i686 unknown"
ALGORITHMPACKAGEACCEPTANCE DATE	"June-1997"
ALGORITHMPACKAGEMATURITYCODE	"at-launch"
ALGORITHMPACKAGENAME #	"ATBD-MOD-02"
ALGORITHMPACKAGEVERSION	"2"
INSTRUMENTNAME	"Moderate Resolution Imaging Spectroradiometer"
LOCALINPUTGRANULEID †	"MOD03.A2001043.1510.005.2005105143551.hdf, MOD021KM.A2001043.1510.005.2005105163716.hdf, MOD35_L2.A2001043.1510.005.2005105171245.hdf"
EXCLUSIONGRINGFLAG	"N"
GRINGPOINTLATITUDE	(-11.9657643150703, -15.1127398445589, -33.2616100115919, -29.5206668804633)
GRINGPOINTLONGITUDE	(-79.8034155352806, -58.1828178475367, -61.0282026524014, -85.5445842551903)
GRINGPOINTSEQUENCENO	(1, 2, 3, 4)
VERYGOODQUALITYDATAPCT_LAND	0.00
GOODQUALITYDATAPCT_LAND	100.00
MARGINALQUALITYDATAPCT_LAND	0.00
BADQUALITYDATAPCT_LAND	0.00
VERYGOODQUALITYDATAPCT_OCEAN	64.65
GOODQUALITYDATAPCT_OCEAN	3.80
MARGINALQUALITYDATAPCT_OCEAN	15.96
BADQUALITYDATAPCT_OCEAN	17.00
ALGORITHMSOFTWAREVERSIONLAND	2
ALGORITHMSOFTWAREVERSIONOCEAN	2
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE04:5.1.0

Refers to Algorithm Theoretical Basis Document (ATBD) reference number

† MODIS product input files

Water Vapor Product: MOD05_L2 (Terra) & MYD05_L2 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Total Precipitable Water Vapor 5-Min L2 Swath 1km and 5km"
PROCESSINGENVIRONMENT	"Linux minion5027 2.4.20-37.7.legacysmp #1 SMP Mon Sep 27 21:38:15 EDT 2004 i686 unknown"
ALGORITHMPACKAGEACCEPTANCEDATE	"June 1997"
ALGORITHMPACKAGEMATURITYCODE	"at-launch"
ALGORITHMPACKAGENAME #	"ATBD-MOD-03"
ALGORITHMPACKAGEVERSION	"2"
INSTRUMENTNAME	"Moderate Resolution Imaging Spectroradiometer"
LOCALINPUTGRANULEID †	"MOD03.A2001043.1510.005.2005105143551.hdf, MOD021KM.A2001043.1510.005.2005105163716.hdf, MOD35_L2.A2001043.1510.005.2005105171245.hdf, MOD07_L2.A2001043.1510.005.2005105171354.hdf"
EXCLUSIONGRINGFLAG	"N"
GRINGPOINTLATITUDE	(-11.9657643150703, -15.1127398445589, -33.2616100115919, -29.5206668804633)
GRINGPOINTLONGITUDE	(-79.8034155352806, -58.1828178475367, -61.0282026524014, -85.5445842551903)
GRINGPOINTSEQUENCENO	(1, 2, 3, 4)
ALGORITHMSOFTWAREVERSION_NIR	2
ALGORITHMSOFTWAREVERSION_IR	2
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE04:5.1.0

Refers to Algorithm Theoretical Basis Document (ATBD) reference number

† MODIS product input files

Cloud Product: MOD06_L2 (Terra) & MYD06_L2 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Clouds 5-Min L2 Swath 1km and 5km"
PROCESSINGENVIRONMENT	"Linux minion5027 2.4.20-37.7.legacysmp #1 SMP Mon Sep 27 21:38:15 EDT 2004 i686 unknown"
ALGORITHMPACKAGEACCEPTANCE DATE	"June-1997"
ALGORITHMPACKAGEMATURITYCODE	"at-launch"
ALGORITHMPACKAGENAME #	"ATBD-MOD-04 and ATBD-MOD-05"
ALGORITHMPACKAGEVERSION	2
INSTRUMENTNAME	"Moderate Resolution Imaging Spectroradiometer"
LOCALINPUTGRANULEID †	"MOD021KM.A2001043.1510.005.2005105163716.hdf, MOD03.A2001043.1510.005.2005105143551.hdf, MOD35_L2.A2001043.1510.005.2005105171245.hdf, Library_water.hdf.v1, Library_ice.hdf.v2, Transmittance.hdf.v2, AlbMap.WS.vb1.0.2001.033.0.659.hdf, AlbMap.WS.vb1.0.2001.033.0.858.hdf, AlbMap.WS.vb1.0.2001.033.1.24.hdf, AlbMap.WS.vb1.0.2001.033.1.64.hdf, AlbMap.WS.vb1.0.2001.033.2.13.hdf, AlbSnwSts.ByNISE.W90.D90.WS.Lats.70.40.2000-2004.Oct.Jan.hdf, IGBP.EcoMap.NtoS.2004.149.v004.hdf"
EXCLUSIONGRINGFLAG	"N"
GRINGPOINTLATITUDE	(-11.9657643150703, -15.1127398445589, -33.2616100115919, -29.5206668804633)
GRINGPOINTLONGITUDE	(-79.8034155352806, -58.1828178475367, -61.0282026524014, -85.5445842551903)
GRINGPOINTSEQUENCENO	(1, 2, 3, 4)
ALGORITHM_VERSION_CLOUD_TOP_PROPERTY_IR	1
ALGORITHM_VERSION_CLOUD_PHASE_IR	1
ALGORITHM_VERSION_CLOUD_PROPERTY_VIS	1
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE06:5.3.0

Refers to Algorithm Theoretical Basis Document (ATBD) reference number

† MODIS product input files

Atmospheric Profile Product: MOD07_L2 (Terra) & MYD07_L2 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Temperature and Water Vapor Profiles 5-Min L2 Swath 5km"
PROCESSINGENVIRONMENT	""Linux minion352 2.4.18-27.7.xsmp #1 SMP Fri Mar 14 05:52:30 EST 2003 i686 unknown""
INSTRUMENTNAME	"Moderate Resolution Imaging Spectroradiometer"
ALGORITHMPACKAGEACCEPTANCE DATE	"June 1997"
ALGORITHMPACKAGEMATURITYCODE	"at-launch"
ALGORITHMPACKAGENAME #	"ATBD-MOD-07"
ALGORITHMPACKAGEVERSION	"2"
LOCALINPUTGRANULEID †	""MOD021KM.A2001043.1510.005.2005105163716.hdf, MOD03.A2001043.1510.005.2005105143551.hdf, MOD35_L2.A2001043.1510.005.2005105171245.hdf, gdas1.PGrbF00.010212.18z, MODIS_REGCOEF_FACTORS.terra.v5, MODIS_senzen.bin, terra_bias.dat.v2, terra_det.dat.v2""
EXCLUSIONGRINGFLAG	"N"
GRINGPOINTLATITUDE	(-11.9657643150703, -15.1127398445589, -33.2616100115919, -29.5206668804633)
GRINGPOINTLONGITUDE	(-79.8034155352806, -58.1828178475367, -61.0282026524014, -85.5445842551903)
GRINGPOINTSEQUENCENO	(1, 2, 3, 4)
PROFILES_ALGORITHM_VERSION_NUMBER	1
STABILITY_INDICES_ALGORITHM_VERSION_NUMBER	1
TOTAL_OZONE_ALGORITHM_VERSION_NUMBER	1
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE03:5.1.5

Refers to Algorithm Theoretical Basis Document (ATBD) reference number

† MODIS product input files

Cloud Mask Product: MOD35_L2 (Terra) & MYD35_L2 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Cloud Mask and Spectral Test Results 5-Min L2 Swath 250m and 1km"
PROCESSINGENVIRONMENT	""Linux minion352 2.4.18-27.7.xsmp #1 SMP Fri Mar 14 05:52:30 EST 2003 i686 unknown""
INSTRUMENTNAME	""Moderate Resolution Imaging Spectroradiometer"
ALGORITHMPACKAGEACCEPTANCE DATE	"June-1997"
ALGORITHMPACKAGEMATURITYCODE	"at-launch"
ALGORITHMPACKAGENAME #	"ATBD-MOD-06"
ALGORITHMPACKAGEVERSION	"2"
LOCALINPUTGRANULEID †	""MOD03.A2001043.1510.005.2005105143551.hdf, MOD021KM.A2001043.1510.005.2005105163716.hdf, MOD02QKM.A2001043.1510.005.2005105163716.hdf, goge1_2_img.v1, ecosystem.img.v1, thresholds.dat.terra.v13, gdas1.PGrbF00.010212.18z, oisst.20010214, eng.010213, NISE_SSMIF13_20010212.HDFEOS"
EXCLUSIONGRINGFLAG	"N"
GRINGPOINTLATITUDE	(-11.9657643150703, -15.1127398445589, -33.2616100115919, -29.5206668804633)
GRINGPOINTLONGITUDE	(-79.8034155352806, -58.1828178475367, -61.0282026524014, -85.5445842551903)
GRINGPOINTSEQUENCENO	(1, 2, 3, 4)
CLOUD_MASK_ALGORITHM_VERSION_NUMBER	1.00
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE03:5.1.5

Refers to Algorithm Theoretical Basis Document (ATBD) reference number

† MODIS product input files

Joint Atmosphere Product: MODATML2 (Terra) & MYDATML2 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Aerosol, Cloud and Water Vapor Subset 5-Min L2 Swath 5km and 10km"
PROCESSINGENVIRONMENT	"Linux minion5027 2.4.20-37.7.legacysmp #1 SMP Mon Sep 27 21:38:15 EDT 2004 i686 unknown"
INSTRUMENTNAME	"Moderate Resolution Imaging Spectroradiometer"
EXCLUSIONGRINGFLAG	"N"
GRINGPOINTLATITUDE	(-11.9657643150703, -15.1127398445589, -33.2616100115919, -29.5206668804633)
GRINGPOINTLONGITUDE	(-79.8034155352806, -58.1828178475367, -61.0282026524014, -85.5445842551903)
GRINGPOINTSEQUENCENO	(1, 2, 3, 4)
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE83:5.1.0

Collection 006 Suggestions:

- 1.) Include a string of PGE's (reflecting the L2 MODIS Atmosphere input products), as was done for L3
(e.g., PRODUCTIONHISTORY = "PGE03:5.1.5; PGE06:5.3.0; PGE04:5.1.0; PGE83:5.1.0")
- 2.) Add LocalInputGranuleID string and populate with the four L2 MODIS Atmosphere input product HDF file names, as is done for all other products. (Note that "InputPointer" in ECS (Core) Inventory Metadata duplicates this info.)

Daily Global Product: MOD08_D3 (Terra) & MYD08_D3 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Aerosol Cloud Water Vapor Ozone Daily L3 Global 1Deg CMG"
PROCESSINGENVIRONMENT	"IRIX64 mtvs2 6.5 10070055 IP35"
LOCALINPUTGRANULEID	†
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE03:5.1.5;PGE06:5.3.0;PGE04:5.1.0;PGE69:5.1.1;PGE56:5.1.1

† MODIS product input files (nearly 1200 L2 MODIS HDF file names)

Eight Day Global Product: MOD08_E3 (Terra) & MYD08_E3 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Aerosol Cloud Water Vapor Ozone Eight Day L3 Global 1Deg CMG"
PROCESSINGENVIRONMENT	"IRIX64 mtvs2 6.5 10070055 IP35"
LOCALINPUTGRANULEID	†
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE03:5.1.5;PGE06:5.3.0;PGE04:5.1.0;PGE69:5.1.1;PGE56:5.1.1;PGE70:5.1.0

† MODIS product input files (up to 8 L3 Daily MODIS HDF file names)

Monthly Global Product: MOD08_M3 (Terra) & MYD08_M3 (Aqua)

Archive Metadata

Field name	Sample Value
LONGNAME	"MODIS/Terra Aerosol Cloud Water Vapor Ozone Monthly L3 Global 1Deg CMG"
PROCESSINGENVIRONMENT	"IRIX64 mtvs2 6.5 10070055 IP35"
LOCALINPUTGRANULEID	†
DESCRREVISION	5.0
PRODUCTIONHISTORY	PGE03:5.1.5;PGE06:5.3.0;PGE04:5.1.0;PGE69:5.1.1;PGE56:5.1.1;PGE57:5.1.0

† MODIS product input files (up to 31 L3 Daily MODIS HDF file names)